

## THE EFFECT OF SODIUM CHLORIDE 0.9% INFUSION SOLUTION AND RINGER'S SOLUTION ON PARTICLE SIZE OF INTRAVENOUS LIPID EMULSION (INTRALIPID 20%)

HAKIM BANGUN\*, JUMIATY, ANAYANTI ARIANTO

Department of Pharmaceutical Technology, Faculty of Pharmacy, Universitas Sumatera Utara, Jl. Tri Dharma No. 5, Pintu 4, Kampus USU Medan 20155, Indonesia. Email: Hakimb17@yahoo.com

Received: 13 September 2016, Revised and Accepted: 14 October 2016

### ABSTRACT

**Objective:** To study the effect of admixing intralipid (IL) with sodium chloride (NaCl) 0.9% infusion solution or Ringer's solution (RS) on physical stability of IL.

**Methods:** Incompatibility test was done in two methods of admixture, i.e. in one container (1:1) and through three-way stopcock (1:2). They were observed for 24 hrs on its emulsion type, pH, microscopic observation of lipid globules, and globules size.

**Results:** The results of admixing IL and electrolyte solution (NaCl or RS) in one container and through three-way stopcock showed no changes in color and emulsion type (O/W). Physical stability evaluation results of IL and NaCl admixture in one container and through three-way stopcock showed that the pH value was 5.1 and 5.2, respectively, and microscopic observation of lipid globules and its size showed coalescence on 24 hrs with size 625.16 nm (one container) and stable during 24 hrs (three-way stopcock). Physical stability evaluation results of IL and RS admixture in one container and through three-way stopcock showed that the pH value was 4.9 and 5.1, respectively, and microscopic observation of lipid globules and its size showed coalescence on 12 hrs with size 572.81 nm (one container) and flocculation on 24 hrs with size 558.12 nm (three-way stopcock).

**Conclusion:** Admixing electrolyte solution (NaCl or RS) and IL in one container and through three-way stopcock causes the larger globule size of IL.

**Keywords:** Intralipid 20%, Sodium chloride 0.9% infusion solution, Ringer's solution, Physical stability.

© 2017 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ajpcr.2017.v10i1.15169>

### INTRODUCTION

Patients in intensive care unit with various and complex conditions receive numerous intravenous (IV) medications. Frequently, the number of IV medication is more than that of venous access lumens for these patients. Therefore, coadministration of IV medications through the same line is done [1]. More than one IV medication increases the risk of incompatibility [2]. However, this situation is poorly understood by health staff [1]. The risks of incompatibility on patients are particulate emboli, tissue irritation, and therapeutic failure [3]. Several researchers have also studied the incompatibility of injection [4,5].

Parenteral nutrition is used when gastrointestinal tracts cannot be accessed and administered depending on patient's condition [6]. Components of parenteral nutrition include amino acid, carbohydrate, lipid, vitamins, trace elements, and electrolyte [7]. Lipid emulsion is important component in parenteral nutrition [8].

Parenteral nutrition solutions are prepared in two forms: 2-in-1 (amino acid and carbohydrate) and 3-in-1 (amino acid, carbohydrate, and lipid) [9]. Advantages of admixing all nutrients in one container provide more convenience for administration, efficient for pharmacy personnel and storage, cheaper, lessen microbe contamination, usage of medical equipment, and monitoring time by nurse [10]. The total parenteral nutrition (3-in-1) that contains lipid tends to cause physical instability with the presence of electrolyte, especially polyvalent cation [11].

Malnutrition continues to be a major health burden in developing countries (for example, Indonesia) and the most important risk factor for illness and death [12,13]. Therapy for malnutrition patients is started with treatment or prevention of hypoglycemia, hypothermia, and dehydration. Hypoglycemia is treated with administration of

dextrose 10% infusion solution [14]. Severe malnutrition is also characterized by essential fatty acid deficiency [13] and is treated by IV lipid emulsion [16]. Electrolyte imbalance is treated by administering Darrow's solution, lactated Ringer's solution (RS), and 0.45% sodium chloride (NaCl) [14].

The previous study by Driscoll *et al.* [17] was done evaluating the stability of amino acid, glucose, and electrolyte solution admixture with various formulations of lipid emulsion 20% which was combination of sunflower oil:soybean oil (1:1) (F1); only soybean oil (F2); mixture of soybean oil:middle chain triglyceride (1:1) (F3); and mixture of olive oil:soybean oil (4:1) (F4). F1 and F2 used egg lecithin as emulsifier, but F3 and F4 used egg lecithin and sodium oleate. Preparation of admixture was done aseptically and it was analyzed in room temperature for 5 times in interval of 30 hrs. The results showed that all formulas were unstable between 8 and 24 hrs, and each admixture showed increment of lipid globules and contained lipid globules which sized 5  $\mu$ m. The research by Cocchio *et al.* [18] was done by admixing IV lipid emulsion with IV medications through Y-site in ratio 1:1 and the results of PFAT5 (proportion of enlarged droplets >5  $\mu$ m) were significantly higher.

Stable IV lipid emulsion contains small droplets with size 200-500 nm. Droplet size  $\geq 5$   $\mu$ m may cause lipid emboli when it enters blood stream [19]. Lipid emboli are occurred on lungs and brain that can cause acute respiratory failure and brain damage, simple anxiety until coma and death [20].

Based on the explanations above, researcher studied the effect of electrolyte solution (NaCl or RS) on physical stability of intralipid (IL) such as alteration in particle size.

## MATERIALS AND METHODS

### Materials

IL (Fresenius Kabi, Jakarta, Indonesia), NaCl (PT. Widatra Bhakti, Pasuruan, Indonesia), RS (PT. Widatra Bhakti, Pasuruan, Indonesia), methylene blue, neutral buffer pH 7.01 (Hanna), acid buffer pH 4.01 (Hanna) and distilled water.

### Methods

#### Admixing in one container

##### Admixing NaCl and IL

NaCl was admixed with IL in ratio 1:1 (12.5:12.5 ml) on room temperature. Then, it was connected with infusion set with flow rate 1 drop/6 seconds, and the admixture result was collected in glass bottle and then its physical stability was evaluated.

##### Admixing RS and IL

RS was admixed with IL in the same ratio and method as stated for admixing NaCl and IL in one container.

#### Admixing through three-way stopcock

##### Admixing NaCl and IL

IL was admixed with NaCl through three-way stopcock in ratio 1:2 (12.5:25 ml) on room temperature with NaCl flow rate was 1 drop/second while IL flow rate was 1 drop/6 seconds. Both infusion sets were connected to three-way stopcock; the admixture result was collected in glass bottle and then its physical stability was evaluated.

##### Admixing RS and IL

IL was admixed with RS through three-way stopcock with the same ratio and method as stated admixing NaCl and IL through three-way stopcock. RS flow rate was 2 drops/second.

#### Physical stability evaluation of NaCl or RS and IL admixture in one container or through three-way stopcock

##### Emulsion type evaluation

A drop of emulsion from admixture result was placed on object glass and then a drop of methylene blue was added. After that, the admixture was homogenized and closed with another object glass.

##### pH determination

pH of admixture was determined using pH meter.

#### Visual observation of lipid globules

Physical appearance evaluation was done within 0, 20, 40, 60 minutes, 3, 6, 12 and 24 hrs with the help of a magnifying glass.

#### Microscopic observation of lipid globules

Lipid globules morphology evaluation was done using digital microscope within 0, 20, 40, 60 minutes, 3, 6, 12 and 24 hrs.

#### Lipid globules size observation

Globules size evaluation was examined using particle size analyzer (Vasco<sup>o</sup>) within 0, 12, and 24 hrs.

## RESULTS

#### Admixing IL and electrolyte solution (NaCl or RS) in one container

IL is a white color emulsion. The results of admixing IL and electrolyte solution (NaCl or RS) in one container showed no changes in color.

#### Admixing IL and electrolyte solution (NaCl or RS) through three-way stopcock

The results of admixing IL and electrolyte solution (NaCl or RS) through three-way stopcock was emulsion with white color too.

#### Physical stability evaluation of NaCl or RS and IL admixture in one container or through three-way stopcock

##### Emulsion type evaluation

As shown in Table 1, it can be observed that admixtures showed no emulsion phase changes during observation time.

##### pH determination

As shown in Table 2, it can be observed that pH of IL reduced when it was admixed with electrolyte solution. One container admixture and calcium-containing admixture (admixture with RS) had lower pH than other admixtures.

#### Visual and microscopic observation of lipid globules

Observation of lipid globules microscopically was done after admixture re-dispersed. Fig. 1a shows no enlargement of lipid globules, and it can be concluded that admixture is stable. Admixtures that showed physical instability (coalescence) are shown in Fig. 1b-d, which were marked by enlargement of lipid globules. Fig. 1e shows flocculation.

#### Lipid globules size observation

As shown in Table 3, it was shown that IL and admixtures of IL with electrolyte solution (NaCl or RS) have lipid droplet size >400 nm which is included in conventional emulsion [34]. From Table 3, it was shown that IL+NaCl in one container 24 hrs, IL+RS in one container 12 and 24 hrs, and IL+RS through three-way stopcock 12 and 24 hrs showed instability due to lipid globules size >500 nm [27].

As shown in Table 3, it can be observed that one container admixtures showed larger lipid globules size than that of admixing through three-way stopcock and also lipid globules size increased on admixture that contained RS. The longer the time after admixing, the larger the lipid globules size. This is caused by due to the longer contact time of positive ion with negative charge of phospholipid so that more lipid globules will unite become larger globules.

Lipid globules' size distribution of IL and admixtures of IL and electrolyte solution (NaCl or RS) on 24 hrs is shown in Fig. 2a-e. The

Table 1: Results of emulsion type evaluation

Time	Emulsion type				
	IL	IL + NaCl (1:1) one container	IL + RS (1:1) one container	IL + NaCl (1:2) three-way stopcock	IL + RS (1:2) three-way stopcock
0 minute	O/W	O/W	O/W	O/W	O/W
20 minutes	O/W	O/W	O/W	O/W	O/W
40 minutes	O/W	O/W	O/W	O/W	O/W
60 minutes	O/W	O/W	O/W	O/W	O/W
3 hrs	O/W	O/W	O/W	O/W	O/W
6 hrs	O/W	O/W	O/W	O/W	O/W
12 hrs	O/W	O/W	O/W	O/W	O/W
24 hrs	O/W	O/W	O/W	O/W	O/W

IL: Intralipid, RS: Ringer's solution, NaCl: Sodium chloride


largest particle distribution is shown in Fig. 2c. which has the longest tail of distribution and the highest intensity peak.

As shown in Table 4, it can be seen that IL+NaCl in one container (24 hrs), IL+RS in one container (12 and 24 hrs), and IL+RS through three-way stopcock (12 and 24 hrs) showed particle size more than 1 µm. IL+RS through three-way stopcock (12 and 24 hrs) showed slower enlargement in size. IL showed a little increment of lipid globules size on 24 hrs due to a little alteration of temperature during observation. The temperature may alterate the size of globules. Nevertheless, the observation was not done in controlled room temperature.


**Table 2: pH of NaCl 0.9%, RS, IL, and its admixtures**

Parenteral solutions	pH
NaCl 0.9%	4.9
RS	5.1
IL	7
IL+NaCl (1:1) 1 container	5.1
IL+RS (1:1) 1 container	4.9
IL+NaCl (1:2) three-way	5.3
IL+RS (1:2) three-way	5.1

IL: Intralipid, RS: Ringer's solution, NaCl: Sodium chloride



**Fig. 1: Microscopic lipid globules observation. (a) Initial; (b) intralipid (IL)+sodium chloride (NaCl) in one container (24 hrs); (c) IL+Ringer's solution (RS) in one container (12 hrs); (d) IL+RS in one container (24 hrs); (e) IL+RS through three-way stopcock (24 hrs)**



**Fig. 2: (a) Lipid globules size distribution of Intralipid (IL) (24 hrs). (b) Lipid globules size distribution of IL which was mixed with sodium chloride (NaCl) in one container (24 hrs). (c) Lipid globules size distribution of IL which was mixed with Ringer's solution (RS) in one container (24 hrs). (d) Lipid globules size distribution of IL which was mixed with NaCl through three-way stopcock (24 hrs). (e) Lipid globules size distribution of IL which was mixed with RS through three-way stopcock (24 hrs)**

**DISCUSSION**

**Emulsion type**

O/W emulsion type is usually given parenterally [22] because blood contains blood cell suspension in plasma solution that has lots of water [21] so that O/W emulsion type is more soluble in blood fluid and more acceptable to the patient.

Physical appearance showed that during 24 hrs, IL which was admixed with electrolyte solution (NaCl or RS) showed coalescence process. This process did not cause phase inversion because the used emulsifier is dissolved in continuous phase (water) which leads to difficulty occurrence of phase inversion for higher volume ratio (water) [23]. Emulsifier in IL is purified phospholipid which the majority contains nonpolar lecithin. However, if lecithin contacts with water, it is hydrated to form emulsion [24]. In other words, the used emulsifier is soluble in water and inversion phase is difficult to occur.

**pH determination**

pH value of IL was obtained 7 while pH value that was stated on product label is ±8 with the adjustment of NaOH aqueous solution. This is caused by when IV lipid emulsion was produced before sterilization, and the pH was adjusted until approximately 8 [25]. Autoclaving process decreases pH and leads to phospholipid hydrolysis to lysophospholipid and free fatty acid so that this process reduces pH to 7 and this value is still acceptable by body [26] and meets the requirements of IV lipid emulsion pH in USP [27]. Slightly alkali condition will cause amphoteric characteristics of lecithin to form negative charge so that the surface charge increases. It also forms thicker bilayer phospholipid films which could form smaller droplet size, increase stability, and had favorable effect in plasma clearance. Na<sup>+</sup> in NaOH formed sodium soaps with free fatty acids and stabilized the submicron emulsion by acting as auxiliary emulsifiers [28].

pH value of other admixtures had lower value than IL and the lowest was shown by RS which was admixed with IL in one container, i.e., 4.9. Lower pH was caused by pH of NaCl and RS itself was low so admixture of IL with NaCl or RS would definitely have lower pH. Lower pH (<5) must be avoided because electrostatic repulsive force among lipid globules lessen and causes globules size enlargement and coalescence [25].

Monovalent cations (Na<sup>+</sup> and K<sup>+</sup>) bind with phosphate groups from phosphatidylcholine and reduce zeta potential of emulsion. Multivalent cation (Ca<sup>2+</sup>) adsorbs hydrophilic parts of phospholipid molecule and reduces zeta potential. Zeta potential becomes smaller with the increasing electrolyte concentration as a result of shrinkage of the electrical double layer at high ionic strength [29]. Based on Debye-Hückel equation, ionic strength of RS (0.158) is higher than NaCl (0.154) [30].

Table 3: Results of admixture lipid globules size

Time (hrs)	Lipid globules size (D mean intensity) (nm)				
	IL	IL+NaCl (1:1) one container	IL+RS (1:1) one container	IL+NaCl (1:2) three-way stopcock	IL+RS (1:2) three-way stopcock
0	447.16	450.73	457.54	445.99	440.56
12	457.33	475.12	572.81	471.51	539.56
24	485.11	625.16	992.37	490.89	558.12

IL: Intralipid, RS: Ringer's solution, NaCl: Sodium chloride

Table 4: Results of admixture lipid globules size range

Time (hrs)	Admixture lipid globules size range (nm)				
	IL	IL+NaCl (1:1) one container	IL+RS (1:1) one container	IL+NaCl (1:2) three-way stopcock	IL+RS (1:2) three-way stopcock
0	186.26-891.49	177.88-891.49	195.04-933.5	195.04-891.49	195.04-891.49
12	204.23-891.49	234.49-891.49	233.49-1230.59	204.23-933.5	223.93-1122.32
24	195.04-1023.56	269.22-1288.59	537.17-1778.75	245.54-891.49	223.93-1175.21

IL: Intralipid, RS: Ringer's solution, NaCl: Sodium chloride

### Visual and microscopic observation of lipid globules

Largest lipid globules microscopically are shown in Fig. 1d due to admixtures of IL and RS in one container. RS contains divalent ion ( $\text{Ca}^{2+}$ ) which is more disruptive than monovalent ions ( $\text{Na}^+$  and  $\text{K}^+$ ) [32]. Electrolyte with divalent charge ( $\text{Ca}^{2+}$ ) bonds strongly with lipid droplet that is stabilized with lecithin to form ion-lipid complex 1:2 so that it forms positive charge [30]. Lipid globules enlargement is caused by coalescence process that can be seen on its physical appearance. Coalescence occurs when lipid droplet was close, thin layer of water between two close droplets ruptures and causes oil from two droplets combine to form larger droplets [33]. Admixtures of IL and NaCl also showed enlargement of lipid globules although it was not as high as admixtures with RS.

IL and electrolyte solution (NaCl or RS) admixture through three-way stopcock achieves more stable condition than admixture in one container as shown in Fig. 1e (flocculation phenomenon) due to the fact that interaction time of both solutions is in shorter time and lots amount of electrolyte solution have been flowed and the remained will interact with lipid globules of IL. Flocculation happens when oil suspended droplet collides due to naturally Brownian motion or agitation. Because of some collisions, attractive force (Van der Waals interaction) occurs between lipid globules because its surface has flaw so that attractive force exceeds repulsion force and causes droplet close to one another [33].

### Lipid globules size observation

Lipid globules size for admixtures that contain RS showed larger globules size. This is due to  $\text{Ca}^{2+}$  in RS which reduces or neutralizes negative charge of phospholipid so that repulsive force decreases between particles and lipid globules coalescence [35].

Human's blood vessel includes aorta, artery, capillary, venule, vena, and vena cava. Smallest human's blood vessel is 5  $\mu\text{m}$ . Therefore, if the size of lipid globules is 5  $\mu\text{m}$  or more, it can cause capillary blockage [36]. Safe IV lipid emulsion should have globules size which is lower than 1  $\mu\text{m}$ . Globules size >1  $\mu\text{m}$  will turn emulsion instable and has potential to enlarge above 5  $\mu\text{m}$  and leads to fat embolism, emboli in capillary and traps on lung capillary so that it affects respiratory function [35,37].

### CONCLUSION

Admixing electrolyte solution (NaCl or RS) and IL in one container or through three-way stopcock causes the larger globule size of IL but admixing in one container causes larger lipid globule size. Furthermore, RS is stronger than NaCl in enlarging lipid globule size.

### ACKNOWLEDGMENT

We would like to thank the Physical Pharmacy Laboratory staffs, Faculty of Pharmacy, University of Sumatera Utara for providing equipment for our study.

### REFERENCES

- Bertsche T, Mayer Y, Stahl R, Hoppe-Tichy T, Encke J, Haefeli WE. Prevention of intravenous drug incompatibilities in an intensive care unit. *Am J Health Syst Pharm* 2008;65(19):1834-40.
- Bergman HD. Incompatibilities in large volume parenterals. *Drug Intell Clin Pharm* 1977;11(6):345-60.
- Melsungen AG. Drug Incompatibility. Germany: B. Braun; 2011. p. 8.
- Konda N, Prashanth P, Arvind G, Shah S. Pharmaceutical development and compatibility studies on cytarabine injection. *Asian J Pharm Clin Res* 2013;6(3):142-5.
- Pimple S, Maurya P, Salunke K, Singh R, Gurjar M, Shah M. Formulation development and compatibility study of dexketoprofen injection used in the management of post-operative pain. *Int J Pharm Sci Rev Res* 2015;30(1):299-305.
- Ansel HC, Prince SJ. *Kalkulasi Farmasetik*. Jakarta: EGC; 2004. p. 163.
- Dietitians Association of Australia. Parenteral Nutrition Manual for Adults in Health Care Facilities. Australia: Dietitians Association of Australia; 2011. p. 9-13.
- Fell GL, Nandivada P, Gura KM, Puder M. Intravenous lipid emulsions in parenteral nutrition. *Adv Nutr* 2015;6(5):600-10.
- Gargasz A. Neonatal and pediatric parenteral nutrition. *AACN Adv Crit Care* 2012;23(4):451-64.
- Krzywda EA, Andris DA, Edmiston CE, Wallace JR. Parenteral access devices. In: ASPEN, editor. *The Science and Practice of Nutrition Support*. USA: American Society for Parenteral and Enteral Nutrition; 2001. p. 257.
- Mirković D, Ibrić S, Antunović M. Quality assessment of total parenteral nutrition admixtures by the use of fractional factorial design. *Vojnosanit Pregl* 2013;70(4):374-9.
- Müller O, Krawinkel M. Malnutrition and health in developing countries. *CMAJ* 2005;173(3):279-86.
- Riskeddas. (2010). *Research Report of Basic Health*. Jakarta: Health Research and Development Administration. Page 23.
- WHO. *Management of Severe Malnutrition: A Manual for Physicians and Other Senior Health Workers*. Geneva: World Health Organization; 1999. p. 1, 7-8.
- Duerksen D, McCurdy K. Essential fatty acid deficiency in a severely malnourished patient receiving parenteral nutrition. *Dig Dis Sci* 2005;50(12):2386-8.
- Calder PC, Waitzberg DL, Koletzko B. *Intravenous Lipid Emulsions*. Germany: S. Karger; 2015. p. 69.
- Driscoll DF, Giampietro K, Wichelhaus DP, Peterss H, Nehne J, Niemann W, et al. Physicochemical stability assessments of lipid

- emulsions of varying oil composition. *Clin Nutr* 2001;20(2):151-7.
18. Cocchio C, Geib AJ, Rynn KO. Physicochemical stability of intravenous fat emulsion in combination with medications used for resuscitation. *SOJ Pharm Pharm Sci* 2014;1(1):1-3.
  19. Schuh RS, Bruxel F, Teixeira HF. Physicochemical properties of lecithin-based nanoemulsions obtained by spontaneous emulsification or high-pressure homogenization. *Quim Nova* 2014;37(7):1193-8.
  20. Filomeno LT, Carelli CR, Da Silvia NC, De Barros Filho TE, Amatuzzi MM. Fat embolism: A review for current orthopaedics practice. *Acta Ortop Bras* 2005;13(4):196-208.
  21. Brust M, Schaefer C, Doerr R, Pan L, Garcia M, Arratia PE, *et al.* Rheology of human blood plasma: Viscoelastic versus newtonian behavior. *American Physical Society* 2013;110(7):1-5.
  22. Mestres GM, Nielloud F. Emulsions in health care applications—An overview. *J Dispers Sci Technol* 2002;23(1-3):419-39.
  23. Vaessen GE, Vissachers M, Stein HN. Predicting catastrophic phase inversion on the basis of droplet coalescence kinetics. *Am Chem Soc* 1996;12(4):875-82.
  24. Rowe RC, Sheskey PJ, Quinn ME. *Handbook of Pharmaceutical Excipients*. 6<sup>th</sup> ed. USA: Pharmaceutical Press; 2009. p. 385, 499, 501.
  25. Floyd AG. Top ten considerations in the development of parenteral emulsions. *PSTT* 1999;2(4):134-43.
  26. Zhao N, Liu X, Sun Y, Wang Y, Pu X, Qin Y, *et al.* Factors affecting particle size of an intravenous fat emulsions. *Asian J Pharm Sci* 2010;5(4):161-7.
  27. USP. *The United States Pharmacopeia*. 36<sup>th</sup> ed. USA: The United States Pharmacopeial Convention; 2012. p. 321-3, 4124-5.
  28. Zhao N, Qin Y, Sun Y, Sun J, He Z. Formulation and preparation of a nitrendipine submicron emulsion. *Asian J Pharm Sci* 2011;6(3-4):151-8.
  29. Wiącek AE. Effect of phospholipid and (phospho) lipase modification on interfacial properties of oil/water emulsion. *UMCS Chem* 2015;52(1):79-109.
  30. Martin AN. *Martin's Physical Pharmacy and Pharmaceutical Sciences*. China: Lippincott Williams & Wilkins; 2011. p. 420-7.
  31. Wermuth CG, Aldous D, Raboisson P, Rognan D. *The Practice of Medicinal Chemistry*. 4<sup>th</sup> ed. USA: Elsevier; 2015. p. 716.
  32. Schroder AM. Total parenteral nutrition - Problems in compatibility and stability. *EJHP Pract* 2008;14(1):65-7.
  33. Baker MT, Naguib M. Propofol: The challenges of formulation. *Anesthesiology* 2005;103(4):860-76.
  34. Burgess DJ. *Injectable Dispersed Systems*. USA: Taylor & Francis Group; 2005. p. 101, 213, 223, 218-9.
  35. Sacks GS. Drug-nutrient considerations in patients receiving parenteral and enteral nutrition. *Pract Gastroenterol* 2004;19:39-48.
  36. Barret KE, Barman SM, Boitano S, Brooks HL. *Ganong's Review of Medical Physiology*. 23<sup>rd</sup> ed. USA: The McGraw-Hill Companies, Inc.; 2010. p. 2.
  37. Wermuth CG, Aldous D, Raboisson P, Rognan D. *The Practice of Medicinal Chemistry*. 4<sup>th</sup> ed. USA: Elsevier; 2015. p. 716.