

THE ROLE OF SELENIUM MICRONUTRIENTS AS ANTIOXIDANTS IN EXPOSURE TO E-CIGARETTE SMOKE

RIVAN VIRLANDO SURYADINATA^{1,2}, MERRYANA ADRIANI³, SANTI MARTINI⁴, SRI SUMARMI³,
BAMBANG WIRJATMADI^{3*}

¹Doctoral Program of Public Health, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia. ²Department of Public Health, Faculty of Medicine, Universitas Surabaya (UBAYA), Surabaya, Indonesia. ³Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia. ⁴Department of Epidemiology, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia
Email: b.wirjatmadi@fkm.unair.ac.id

Received: 09 July 2019, Revised and Accepted: 15 July 2019

ABSTRACT

Objective: E-cigarette products have resulted in various controversies concerning their posed impacts on health. Some argue that exposure to e-cigarette smoke could improve free radicals in the body; thus, it causes harming impacts on health. Peroral selenium (Se) administration can increase superoxide dismutase (SOD) and glutathione peroxidase (GPx) serving as antioxidants in the body.

Methods: This research is an experimental study aiming to analyze the effectiveness of Se to decrease free radical due to exposure to e-cigarette smoke as one of the preventive actions. The research was carried out to male Wistar rats with exposure to e-cigarette smoke and peroral Se intake with different time and duration of administration.

Results: Research results showed a decrease of antioxidant SOD and GPx in the administration of exposure to e-cigarette smoke, and they gradually increased after Se administration ($p=0.000$). Meanwhile, the malondialdehyde level was inversely proportional compared to antioxidant SOD and GPx.

Conclusion: Se is a micronutrient that can reduce free radicals due to exposure to e-cigarette smoke through enhancement of antioxidant enzymes such as SOD and GPx.

Keywords: Selenium, Malondialdehyde, E-cigarette, Glutathione peroxidase, Superoxide dismutase.

© 2019 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ajpcr.2019.v12i8.34454>

INTRODUCTION

The use of electronic cigarettes keeps on increasing annually. It is caused by the absence of specific regulations targeting the use of e-cigarettes leading to their use at any place [1]. Besides, their great shapes and models compared to tobacco cigarettes along with their massive promotion attract young people [2]. Electronic cigarettes are also claimed as tools to reduce or stop the use of tobacco cigarettes. They are also considered to be safer and have less risk because they only produce water vapor claimed not to endanger health [3]. Various studies have shown the dangerous risks posed by exposure to electronic cigarettes. The danger of electronic cigarettes has been estimated to be greater compared to the one in tobacco cigarettes. Even in passive smokers inhaling the smoke of e-cigarettes, nicotine is also found in blood [4]. The inhaled smoke of e-cigarettes entering airways can trigger the existence of free radicals. The amount enhancement of excessive free radicals will cause amount imbalance of free radicals and antioxidants in the body so that oxidative stress occurs [5].

Free radicals in the body are molecules having one or more electrons which are not in pairs, but very reactive, and unstable. Free radicals can react fast with other molecules by catching electrons to stabilize themselves. Free radicals will be balanced by taking electrons, whereas the molecules under attacked will change into other free radicals causing damage to cells [6]. Oxidative stress will enhance lipid peroxidase on the attacked cell membrane [7]. One of lipid peroxidase results due to free radicals is malondialdehyde (MDA), which can be used as a parameter of free radicals enhancement in the body [8].

Reduction of free radicals can be done by administering antioxidants. Antioxidants mechanism works by giving electrons to the free radicals; thereby, they can stop cell damage process [9]. Besides, antioxidants

will neutralize free radicals so that they will not have any ability to take electrons from the cells [10]. Physiologically, human body can neutralize free radicals when their amount is not excessive [11]. This is done by means of endogenous antioxidant defense mechanism [12,13]. However, when there is a lack of endogenous antioxidants, the body needs exogenous antioxidants [14]. Various ways to enhance antioxidants in the body include adding intake of exogenous antioxidants from food, supplements, and doing sports [15]. Some of the endogenous antioxidants needed to neutralize free radicals free are glutathione peroxidase (GPx) and superoxide dismutase (SOD). Antioxidant GPx is an intracellular antioxidant enzyme which enzymatically changes hydrogen peroxidase (H_2O_2) to water (H_2O) [16]. Meanwhile, antioxidant SOD plays a role in changing superoxide radical into H_2O_2 [17].

Selenium (Se) has antioxidant and immunomodulation functions. In the body, Se is used to synthesize amino acid, called selenocysteine, which is very important for selenoprotein formation. Selenoprotein is composed of 25 types in human body and classified based on its function as various enzymes which one of them is antioxidant GPx [18]. Thus, Se is often called as an enzymatic component of GPx [19]. The main form of Se is selenomethionine, commonly found in food [20]. Main sources of Se in food are selenomethionine amino acid (in the form of cereals) and selenocysteine (from animal products). Some plants such as beet leaves, cabbage, and garlic contain a high level of Se, reaching up to 50% in the form of selenate. Se can also be obtained from supplements in the forms of selenate, selenite, selenomethionine, or other forms of Se [21]. Moreover, Se has a quite significant function for human's health because it is able to join into protein to be selenocysteine. Se effects have been proven, ranging from antioxidant to anti-inflammatory effects [22].

Various studies have proven that SE suppresses reactive oxygen species formation by improving GPx activity [23]. Antioxidant SOD is more influenced by copper and zinc micronutrients. Hence, the role of Se against SOD enhancement still needs further research [24].

METHODS

This research is an experimental study using a post-test control group design (Ethical Approval No:103/EA/KEPK/2018). The research was conducted for 5 weeks utilizing male Wistar rats (*Rattus norvegicus*). The research was divided into six groups with different treatments in each group.

The first group was the negative control group where the experimental animals did not undergo any intervention for 4 weeks. The second group was the cigarette control group where the experimental animals were given intervention of exposure to e-cigarette smoke for 4 weeks. The third group was the Se control group where the experimental animals were given intervention of peroral Se intake for 4 weeks. The fourth group was treatment Group I where the experimental animals were given exposure to e-cigarette smoke on the 1st week. Next, on the second until the 5th week, they were given exposure to e-cigarette smoke and peroral Se. The fifth group was treatment Group II, where the experimental animals were given exposure to e-cigarette smoke and peroral Se for 4 weeks. The sixth group was treatment Group III where the experimental animals were given peroral Se in the 1st week, next during the second until the 5th week they were given exposure to e-cigarette smoke and peroral Se.

Mice

Male Wistar rats aged 2–3 months weighing 150–200 g were used as animal models. The Wistar rats were never been subjects research and did not have macroscopic abnormalities. The experimental animals are adapted first for 5 days before giving intervention. The research was conducted in the Laboratory of the Faculty of Medicine, Universitas Airlangga.

Se

Se per oral used is an active form of selenomethionine. The recommended dose for adults is 200 mcg/day. Using the conversion table Laurence and Bacharach, calculation obtained was 3.6 mcg. Thus, the dosage that can be given to male white rats of the Wistar strain is 3.6 mcg/day per each experimental animal.

E-cigarette

The solution of an e-cigarette used contains 6 mg of nicotine. The room where exposure to e-cigarette smoke measures 50 cm × 40 cm × 20 cm there is space a pipe will be passed which can expend e-cigarette smoke. The exposure to e-cigarette smoke exposure is adjusted for the length of time planned for the research.

Measurement of GPx level in the blood

GPx level in blood measured using an enzyme-linked immunosorbent assay (ELISA) Kit, which can be applied in the quantitative *in vitro* determination of Rat GPx1 concentration in serum, plasma, and other biological fluids. The ELISA kit uses the competitive-ELISA method. The kit had been coated with Rat GPx1 as a specific antibody. The rat GPx1 concentration in the sample was then determined by comparing the sample result to the standard curve.

Measurement of SOD level in blood

SOD level in blood measured using an ELISA kit which can be applied in the quantitative *in vitro* determination of rat SOD1 concentration in serum, plasma, and other biological fluids. The ELISA kit uses the competitive-ELISA method. The kit had been coated with rat SOD1 as a specific antibody. The rat SOD 1 concentration in the sample was then determined by comparing the sample OD to the standard curve.

Measurement of MDA level in blood

MDA levels in blood measured using thiobarbituric acid (TBA)

reactive substance (TBARS) assay. Measurements were carried out using the bioassay system based on the reaction of TBARS with TBA in forming pink-colored compounds. The color intensity was measured at 535 nm or with fluorescence intensity at 560 nm/585 nm which was directly proportional to the concentration of TBARS in the sample.

Statistika

The collected data will be statistically tested using SPSS version 20 to analyze the difference and influence of Se on SOD levels, GPx, and MDA in the blood in all groups.

RESULTS

SOD levels in blood as a result of exposure to e-cigarette smoke

Research results were obtained by comparing the levels of SOD in each group. Based on Fig. 1, the average value and difference of levels of SOD in each group can be known. The results showed that the cigarette control group had the lowest average value reaching 2.29±0.12. Meanwhile, in the Se control group, it had the highest average value reaching 5.05±0.16. In the treatment groups, there was an increase in the average value of SOD sequentially.

To fulfill the parametric test requirements, the research results were subjected to a normality test ($p>0.05$) and homogeneity test ($p=0.008$). Based on Table 1, Kruskal-Wallis test was conducted to know the difference of SOD levels in the groups ($p=0.000$).

GPx levels in blood as a result of exposure to e-cigarette smoke

Research results were obtained by comparing levels of GPx in each group. Based on Fig. 2, the average value and difference of levels of GPx in each group can be known. The results showed that the cigarette control group had the lowest average value reaching 85.6±8.96. Meanwhile, in the Se control group, it had the highest average value reaching 330.6±13.22. In the treatment groups, there was an increase in the average value of GPx sequentially.

To fulfill the parametric test requirements, the research results were subjected to normality test ($p>0.05$) and homogeneity test ($p=0.385$). Based on Table 2, analysis of variance test was conducted to know the difference of GPx levels in the groups ($p=0.000$).

Table 1: Kruskal-Wallis test for superoxide dismutase in each group

Groups	Definition	Mean±SD	p-value
I	Negative control group	4.64±0.27	0.000
II	Cigarette control group	2.29±0.12	
III	Selenium control group	5.05±0.16	
IV	Treatment Group I	3.00±0.04	
V	Treatment Group II	3.33±0.1	
VI	Treatment Group III	4.67±0.07	




Fig. 1: The mean value for superoxide dismutase in each group




Fig. 2: The mean value for glutathione peroxidase in each group




Fig. 3: The mean value for malondialdehyde in each group

MDA levels in blood as a result of exposure to e-cigarette smoke

Research results were obtained by comparing levels of MDA in each group. Based on Fig. 3, the average value and difference of levels of MDA in each group can be known. The results showed that the negative control group had the lowest average value reaching 0.0144 ± 0.0018 . Meanwhile, in the cigarette control group, it had the highest average value reaching 0.2416 ± 0.0174 . In the treatment groups, there was an decrease in the average value of MDA sequentially.

To fulfill the parametric test requirements, the research results were subjected to normality test ($p > 0.05$) and homogeneity test ($p = 0.000$). Based on Table 3, Kruskal-Wallis test was conducted to know the difference of MDA levels in the groups ($p = 0.000$).

DISCUSSION

E-cigarette smoke entering the airways will increase free radicals in the body. The enhancement of free radicals can trigger an increase of MDA as well as a decrease of the body's enzymatic antioxidants in the forms of SOD and GPx [25]. The free radicals are also actually needed in the body because they play an important role in the body's defense system [26]. However, the imbalance between amount of free radicals and antioxidants will cause the occurrence of cell damage [27,28]. Additional intake of selenium per oral is needed to stimulate antioxidant increase in the body [29,30].

Administering Se in this research showed enhancement existence of SOD and GPx. The group with the highest increase of antioxidants was the Se control group. It was proven by higher enhancement of antioxidants compared to the negative control groups. Meanwhile, the cigarette control group had the lowest level of antioxidants. This was caused by an increase of the amount of free radicals entering the airways because of exposure to e-cigarette smoke. On the three treatment groups, there was antioxidant enhancement gradually. The first treatment group had a higher level of antioxidant due to the administration of Se done earlier than exposure to e-cigarette smoke. As a result, the body's enzymatic

Table 2: Analysis of variance test for glutathione peroxidase in each group

Groups	Definition	Mean \pm SD	p-value
I	Negative control group	236.2 \pm 7.3	0.000
II	Cigarette control group	85.6 \pm 8.96	
III	Selenium control group	330.6 \pm 13.22	
IV	Treatment Group I	118.2 \pm 6.46	
V	Treatment Group II	154.0 \pm 5.66	
VI	Treatment Group III	256.4 \pm 6.69	

Table 3: Kruskal-Wallis test for malondialdehyde in each group

Groups	Definition	Mean \pm SD	p-value
I	Negative control group	0.0144 \pm 0.0018	0.000
II	Cigarette control group	0.2416 \pm 0.0174	
III	Selenium control group	0.0204 \pm 0.0028	
IV	Treatment Group I	0.2104 \pm 0.0051	
V	Treatment Group II	0.1802 \pm 0.0043	
VI	Treatment Group III	0.1410 \pm 0.0032	

antioxidants were still able to reduce the free radicals entering through the airways. In the second treatment group, the administration of both exposures to e-cigarette and Se was done at the same time. Therefore, there was a decrease in the antioxidant level. The third treatment group indicated the lowest level of antioxidants of all other treatment groups. It happened because the free radicals had entered the airways earlier. Hence, the antioxidant enhancement did not show any significant increase.

Se is a supplement that plays an important role in human's health. The given Se intake is expected to be able to prevent the increase of antioxidants SOD and GPx as well as provide with potential hope as a therapy to prevent cell damage caused by exposure to e-cigarette [31].

CONCLUSION

Se is a micronutrient that can reduce free radicals due to exposure to e-cigarette smoke through enhancement of antioxidant enzymes such as SOD and GPx.

AUTHORS' CONTRIBUTIONS

All the authors have contributed equally in research, review, and finalization of the contents of the journal.

CONFLICTS OF INTEREST

Declare None.

REFERENCES

- Canistro D, Vivarelli F, Cirillo S, Marquillas CB, Buschini A, Lazzaretti M, et al. E-cigarettes induce toxicological effects that can raise the cancer risk. *Sci Rep* 2017;7:2028.
- Grana R, Benowitz N, Glantz SA. E-cigarettes: A scientific review. *Circulation* 2014;129:1972-86.
- Vardavas CI, Anagnostopoulos N, Kougias M, Evangelopoulou V, Connolly GN, Behrakis PK, et al. Short-term pulmonary effects of using an electronic cigarette: Impact on respiratory flow resistance, impedance, and exhaled nitric oxide. *Chest* 2012;141:1400-6.
- Polosa R, Caponnetto P, Morjaria JB, Papale G, Campagna D, Russo C, et al. Effect of an electronic nicotine delivery device (e-cigarette) on smoking reduction and cessation: A prospective 6-month pilot study. *BMC Public Health* 2011;11:786.
- Suryadinata RV, Wirjatmadi B, Adriani M. Effectiveness Decrease Combined with Supplements Malondialdehyde Antioxidant Superoxide Dismutase Gliadin Melon with Due to Exposure to Cigarette. *Glob Med Heal Commun* 2017;5:79-83.
- Suryadinata RV. Effect of free radicals on inflammatory process in chronic obstructive pulmonary disease (COPD). *Amerta Nutr*

- 2018;2:317-24.
7. Raghavendra U, Rao A, D'souza J, Pai VR, Nair S, Kumar V, et al. Comparative estimation of salivary total antioxidant capacity in periodontal health and chronic periodontitis a pilot study. *Asian J Pharm Clin Res* 2018;11:523.
 8. Ayala A, Muñoz MF, Argüelles S. Lipid peroxidation: Production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid Med Cell Longev* 2014;2014:360438.
 9. Mapunda PE, Mligo C, Lyaruu HV. Evaluation of free radical scavenging ability and antiradical activities of *Ximenia caffra* fruit extracts at different ripening stages. *Int J Pharm Pharm Sci* 2019;11:55.
 10. Pratiwi SR, Lorensia A, Suryadinata RV. Asupan Vitamin C dan E dengan SQ-FFQ terhadap fungsi paru perokok dan non-perokok. *Media Kesehatan Masyarakat Indones* 2018;14:101.
 11. Subba A, Mandal P. Standardization of a traditional polyherbal formulation with pharmacognostic study; its phytochemical content, antioxidant, and antidiabetic activity. *Asian J Pharm Clin Res* 2019;12:5.
 12. Rahal A, Kumar A, Singh V, Yadav B, Tiwari R, Chakraborty S, et al. Oxidative stress, prooxidants, and antioxidants: The interplay. *Biomed Res Int* 2014;2014:761264.
 13. Pratima H. Antioxidant and antibacterial activity of alkaloid extract of *Cucumis trigonus* Roxb. *Int J Pharm Pharm Sci* 2019;11:44-8.
 14. Liguori I, Russo G, Curcio F, Bulli G, Aran L, Della-Morte D, et al. Oxidative stress, aging, and diseases. *Clin Interv Aging* 2018;13:757-72.
 15. Suryadinata RV, Wirjatmadi B, Adriani M. Effects of Changes in Goblet Cell Hyperplasia for 28 Days of Cigarette Smoke Exposure through the administration of Antioxidant Superoxide Dismutase. *Indones J Public Health* 2017;11:60.
 16. Lubos E, Loscalzo J, Handy DE. Glutathione peroxidase-1 in health and disease: From molecular mechanisms to therapeutic opportunities. *Antioxid Redox Signal* 2011;15:1957-97.
 17. Phaniendra A, Jestadi DB, Periyasamy L. Free radicals: Properties, sources, targets, and their implication in various diseases. *Indian J Clin Biochem* 2015;30:11-26.
 18. Kim Y, Kim DC, Cho ES, Ko SO, Kwon WY, Suh GJ, et al. Antioxidant and anti-inflammatory effects of selenium in oral buccal mucosa and small intestinal mucosa during intestinal ischemia-reperfusion injury. *J Inflamm (Lond)* 2014;11:36.
 19. Socha K, Kochanowicz J, Karpińska E, Soroczyńska J, Jakoniuk M, Mariak Z, et al. Dietary habits and selenium, glutathione peroxidase and total antioxidant status in the serum of patients with relapsing-remitting multiple sclerosis. *Nutr J* 2014;13:62.
 20. Bermingham EN, Hesketh JE, Sinclair BR, Koolaard JP, Roy NC. Selenium-enriched foods are more effective at increasing glutathione peroxidase (GPx) activity compared with selenomethionine: A meta-analysis. *Nutrients* 2014;6:4002-31.
 21. Mehdi Y, Hornick JL, Istasse L, Dufrasne I. Selenium in the environment, metabolism and involvement in body functions. *Molecules* 2013;18:3292-311.
 22. Cai X, Wang C, Yu W, Fan W, Wang S, Shen N, et al. Selenium exposure and cancer risk: An updated meta-analysis and meta-regression. *Sci Rep* 2016;6:19213.
 23. Kim KS, Suh GJ, Kwon WY, Kwak YH, Lee K, Lee HJ, et al. Antioxidant effects of selenium on lung injury in paraquat intoxicated rats. *Clin Toxicol (Phila)* 2012;50:749-53.
 24. Younus H. Therapeutic potentials of superoxide dismutase. *Int J Health Sci (Qassim)* 2018;12:88-93.
 25. Suryadinata RV, Wirjatmadi B, Adriani M, Sumarmi S. The effects of exposure duration to electronic cigarette smoke on differences in superoxide dismutase and malondialdehyde in blood of wistar rats. *Int J Curr Pharm Res* 2019;11:13-6.
 26. Chidambararajan P, Keerthana V, Priyadharshini K, Sakthivel B. *In vitro* antioxidant and anticancer activity of ulva lactuca l. Using MOLT-3 cell line. *Asian J Pharm Clin Res* 2019;12:3-6.
 27. Simioni C, Zauli G, Martelli AM, Vitale M, Sacchetti G, Gonelli A, et al. Oxidative stress: Role of physical exercise and antioxidant nutraceuticals in adulthood and aging. *Oncotarget* 2018;9:17181-98.
 28. Jing SL, Yen KP, Dash GK. *In vitro* antioxidant and photoprotective activities of *Carica papaya* fruits. *Asian J Pharm Clin Res* 2019;12:2-4.
 29. Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn Rev* 2010;4:118-26.
 30. Pizzino G, Irrera N, Cucinotta M, Pallio G, Mannino F, Arcoraci V, et al. Oxidative stress: Harms and benefits for human health. *Oxid Med Cell Longev* 2017;2017:8416763.
 31. Wang N, Tan HY, Li S, Xu Y, Guo W, Feng Y, et al. Supplementation of micronutrient selenium in metabolic diseases: Its role as an antioxidant. *Oxid Med Cell Longev* 2017;2017:7478523.