

## EFFECTIVENESS OF A HEALTH PROMOTION PROGRAM USING THE INTERNATIONAL ORGANIZATION FOR STANDARDIZATION IN KLONGYONG AND NIKHOMPATTANA, THAILAND

SUCHINDA JARUPAT<sup>1</sup> MARUO ,NOPPAWAN PIASUE<sup>2</sup>, SURINTORN KALAMPAKORN<sup>1</sup>, PINYO UTTHIYA<sup>2</sup>, SIRIPORN SASIMONTHONKUL<sup>3</sup>, HISA AKI TABUCHI<sup>4</sup>, HISAKI AKASAKI<sup>5\*</sup>, TOSHI AKI WATANABE<sup>6</sup>, NAOYA TAKI<sup>6</sup>, RYOJI UCHIYAMA<sup>7</sup>, KAZUKI ASHIDA<sup>7</sup>, MASAO OKUHARA<sup>8</sup>, KOJI TERASAWA<sup>6</sup>

<sup>1</sup>Department of Occupational Health and Safety, Faculty of Public Health, Mahidol University, 420/1 Ratchathewi, Bangkok 10400, Thailand. <sup>2</sup>Department of Nursing, Ramathibodi School of Nursing, Faculty of Medicine, Ramathibodi Hospital, Mahidol University, 270 Rama IV Road, Ratchathewi, Bangkok 10400, Thailand. <sup>3</sup>Department of Rehabilitation and Movement Sciences, Faculty of Sports Science Kasetsart University, Kamphaeng Saen Campus, 1 Moo 6. Malaiman Rd., Kamphaeng Saen, Nakonpathom 73140, Thailand. <sup>4</sup>Department of Psychology, University of Innsbruck, Innrain 52, 6020 Innsbruck, Austria. <sup>5</sup>Academic Assembly School of Science and Technology Institute of Engineering, Wakasato, Naganoshi, Nagano 380-0928, Japan. <sup>6</sup>Department of Sports Sciences, Faculty of Education, Shinshu University, 6-Ro Nishinagano, Nagano City, Nagano 380-8544, Japan. <sup>7</sup>Department of Electronics and Computer Science, National Institute of Technology, Nagano College, 716 Tokuma, Nagano City, Nagano 381-8550, Japan. <sup>8</sup>Department of Applied Information Engineering, Faculty of Engineering, Suwa University of Science, 5000-1 Toyohara, Chino City, Nagano 391-0292, Japan.

Email: kterasa@shinshu-u.ac.jp

Received: 29 November 2020, Revised and Accepted: 24 February 2020

### ABSTRACT

**Objectives:** This study aimed to establish a health-care program that adopted ISO9001:2008 (the International Organization for Standardization) in Thailand to improve problems and inspect its effectiveness. Furthermore, we make this health promotion of ISO widely available in Asian countries and make an international contribution.

**Methods:** We implemented a 9-month health program in the Klonyong, and a 6-month health program in Nihompattana, Rayong, Thailand. This program assessed findings from pedometer, anthropometry, physical fitness, and brain function tests.

**Results:** In Klonyong, the average number of walking and exercise steps was 3471.3±129.1, and in Nihompattana, the average number of walking and exercise steps was 4695.5±408.3. The pre- and post-health programs in Klonyong showed significant differences in blood pressure, hand grip strength, 10 m-obstacle walk, and 6-min walk, and in Nihompattana, significant differences in the hand grip strength, sit and reach flexibility, and brain function test were observed. The pre- and post-health programs in Klonyong and Nihompattana showed significant differences in a total number of miss times.

**Conclusions:** The findings from the before and after pre-and post-health programs in Nihompattana suggest that the increased physical activity during the course of the program may have led to improved brain function results.

**Keywords:** Health promotion, Pedometer, Physical fitness, Brain function.

© 2020 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ajpcr.2020.v13i3.36506>

### INTRODUCTION

The World Health Organization has set the goal of "Primary Health Care" (PHC). PHC recognizes all people's health as a basic human right and guarantees subjective participation and self-determination rights in the process of achieving it [1]. This concept led to the health promotion of 1986 [2]. Health promotion is an important tool in preventing illness [3,4]. The Chuan Government's decentralization policy was instrumentally used by community health-oriented bureaucrats in pursuing legislation or the National Health Act, considered a concrete expression of the health promotion movement [5].

The number of elderly people around the world is expected to increase. In Thailand, the number of the elderly people aged over 65 years is increasing. The United Nations (UN) reported that an "aging society" is when >7% of the total population is 65 years old or over, and an "aged society" is when that number increases to >14%. The UN reported that Singapore and Thailand, countries of the Association of Southeast Asian Nations, are aging fast and the population 65 years old and over is projected to increase by more than 7 million in Singapore and more than 1 million in Thailand, from 2020 to 2025 [6].

It is important for health care to provide information, to share it, to cooperate with everyone, and to clarify the role sharing of counselors and coaches [7].

To achieve the elderly population's desired health and well-being, a thorough assessment of health and health behaviors would be the foundation for tailoring an effective plan. In social ecological models, interactions between community members and their environment are crucial. Assessments of the health and health behaviors are the foundation for tailoring a health promotion-prevention plan to a given client. The assessment provides the database for making clinical judgments regarding the client's health strengths, health problems, nursing diagnoses, and desired health or behavioral outcomes, as well as the interventions likely to be effective [8].

The health education curriculum in Thailand focuses on nutrition and exercise [9,10]. The health education curriculum developed for this study, which adopted ISO9001:2008 (the International Organization for Standardization), has been implemented by measuring energy expenditure and conducting brain function, physical fitness, and blood tests and providing educational seminars regarding exercise

and nutrition and recreational activities, such as hiking and cooking [11-14]. The practice of health education in Thailand usually includes monthly meetings with club members and periodic seminars to develop participants' leadership through Thai yoga and stretch exercises; however, the absence of a systematic evaluation after initiation of the program was observed to make an evaluation of the effectiveness of such a health program difficult for the Thai health practitioners. This study aimed to establish our health-care program in Thailand by adopting ISO9001:2008, improving on its problems, and using its effectiveness to compare the health education of Klonyong and Nihompattana in Thailand.

## METHODS

### General method

We implemented a 9-month health program from December 2014 to August 2015 in Klonyong, and a 6-month health program from August 2013 to January 2014 in the city of Nihompattana, in Thailand. This program assessed findings from pedometry, anthropometry, physical fitness, and brain function tests. For the measurement and testing of this study, we first performed a blood test and then measured weight, height, and blood pressure. Second, we had participants perform the go/no-go task, which is a brain function measurement, and third, we performed physical fitness measurement by measuring hand grip strength, sit-ups, sit and reach flexibility, an eyes-open single-leg stance, and a 10-m obstacle walk. Finally, we measured the distance of a 6-min walk.

The study participants from Klonyong were 46 elderly participants aged 69.8±6.5 years, including 13 men aged 69.9±6.4 years and 33 women aged 69.7±6.7 years. The study participants from Nihompattana were 114 elderly participants aged 60.5±11.1 years, including 31 men aged 66.3±7.1 years and 83 women aged 58.3±11.5 years. The participants were healthy elderly people from Klonyong and Nihompattana, Thailand, who attended public health center program once a month and agreed to be participants in this study. Those who lost their health during this health education implementation period and were unable to come to the health center were excluded from the study.

In Klonyong and Nihompattana, during this program, the participants received a series of seminars regarding recreational activities for 90 or 120 min once or twice per month. In addition, the subjects from Klonyong performed stretching and muscular training for 90 min once per week, and Nihompattana performed aerobic exercises 3 times per week and at-home training every day (Table 1). Because the results of the brain function, physical fitness, body weight, and blood chemistry tests' values were significantly improved by the exercise of 7000 or more steps [11-14]. The latest guidelines based on the Helsinki Declaration were adopted by the Institutional Ethics Committee of Mahidol University (Mahidol Univ. ID: 01-58-10) and Shinshu University (UMIN00009309). Written informed consent was obtained from all the participants.

### Pedometry

A pedometer measured the daily number of walking steps and amount of energy expenditure (Acos Co. Ltd., Japan; AM500NE) to count the daily and exercise steps. Exercise steps are defined as steps taken during expenditure >4 METS. The pedometer enabled the data to be transferred and saved to a personal computer. These walking steps were measured from December 2014 to August 2015 in Klonyong and from August 2013 to January 2014 in Nihompattana. The participants reported their results to a project leader during a monthly meeting.

### Anthropometry and blood pressure measurements

The anthropometric measurement adopted weight and body mass index (BMI) measurements. The weight measurement used body composition monitors with scales (Omron Healthcare Co., Ltd. JAPAN; HBF-359). Maximum and minimum blood pressures were measured using auscultation (mercury sphygmomanometer; Kenzumedico 0601B001, Japan), after the study participants had been sitting for

**Table 1: Program contents of the health education in the Klonyong and Nihompattana groups**

Month	Klonyong program	Nihompattana program
August	Measurement before the health education	Measurement before the health education
September		Aerobic exercises three times a week
October		Aerobic exercises three times a week
November		Lecture on blood pressure and aerobic exercises three times a week
December	Measurement before the health education	Visit of the temple and aerobic exercises three times a week
January	Lecture of the nutrition, stick exercise, brain train	Measurement after completion of the program
February	Lecture of the nutrition, stick exercise, recreation	
March	Lecture of the hypertension, Tai Chi Chuan, recreation	
April	Lecture of the nutrition, stick exercise, brain training	
May	Lecture of the diabetes, stick exercise, aroma therapy	
Jun	Lecture of the nutrition, aroma aerobics exercises	
July	Lecture of the hyper cholesterol, stick e brain training	

Klonyong group: Monthly classroom activity plus weekly 90 min strength and weight training, Nihompattana group: Monthly classroom activity plus aerobic exercises 3 times per week and at-home training every day

15 min in a room with an ambient temperature of 25°C and relative humidity of approximately 50%.

### Physical fitness tests

The physical fitness tests administered in this study were approved by the Japanese Ministry of Education, Culture, Sports, Science, and Technology [15]. The physical fitness test (target age: 65–79 years) included six physical assessments: (1) Grip strength for muscle strength; (2) sit-ups for muscle endurance; (3) sit and reach flexibility for muscle flexibility; (4) eyes-open single-leg stance for balance ability; (5) 10-m obstacle walk for walking ability; and (6) a 6-min walk for endurance. The study participants' physical ability was assessed before and after the health program.

### Brain function tests

The go/no-go task [16-18] was used to estimate the inhibitory decision process and comprised three experimental stages: (1) Formation; (2) differentiation, and (3) reverse differentiation. First, in the formation stage, participants were instructed to squeeze a rubber ball in response to a red light that was randomly displayed. The formation stage comprised five trials. Second, during the differentiation stage, the participants squeezed a rubber ball in response to a red light, but not a yellow light, when a red or yellow light was randomly displayed. Third, during the reverse differentiation stages, the participants squeezed a rubber ball in response to yellow light, but not a red light, when a red or yellow light was randomly displayed. In each of the differentiation and reverse differentiation stages, the participants completed 20 trials. Red and yellow lights were equally randomly displayed 10 times each. In this article, the term "miss" indicates an incorrect response when the participants did not squeeze a rubber ball when it should have been squeezed.

Conversely, the term "mistake" means an incorrect response when the participants squeezed the rubber ball when it was not supposed to be squeezed. The participants go/no-go task was assessed before and after the health program.

### Statistical analysis

The nonpaired t-test was used to compare the results before and after participation for the steps measured by the pedometer. The paired t-test was used to compare the results before and after participation in the health program. A 2×2 (before and after for each group: Klongyong and Nihompattana) two-way analysis of variance (ANOVA) with repeated measures on the variable was performed to evaluate for significance in the scores of anthropometry, blood pressure, physical fitness, and go/no-go task. Following the significant interaction, a one-way ANOVA was calculated to interpret the before and after results for Klongyong and Nihompattana. *Post hoc* tests were performed using Tukey–Kramer correction. The level of significance was set at  $p < 0.05$ . Statistical analyses were performed using the SPSS 11.0.1 statistical package (SPSS Inc., Chicago, USA).

## RESULTS

### Pedometry

Fig. 1 shows the average daily number of walking and exercise steps for each month in Klongyong and Nihompattana. In Klongyong, the monthly steps were as follows: December; 3973.6±120.2 (mean±standard error) and exercise steps; 1909.5±137.2, January; 3804.3±139.4 and 1856.2±162.2, February; 4162.4±179.0 and 1631.5±164.9, March; 4070.8±149.9 and 1725.0±106.6, April; 3090.7±104.5 and 1218.4±68.3, May; 2744.0±74.9 and 941.9±45.8, June 2795.2±74.7; 869.8±42.7, July; 3314.8±123.3 and 1166.0±70.0, August; 3285.9±195.9 and 1155.7±121.6. The average number of steps decreased from April to August.

For the 9-month health program, the average walking steps were 3741.3±129.1; the average exercise steps were 1386.0±102.2. In Nihompattana, the monthly steps observed are as follows: August walking steps 4216.2±440.8 and exercise steps 1502.0±568.3; September, 5031.2±366.8 and 1501.0±127.7; October, 4841.5±314.9 and 1452.9±98.7; November, 4670.1±303.2 and 1317.5±86.2; December, 4751.8±374.3 and 1294.3±112.3; and January, 4662.3±396.5 and 1301.3±112.1. The average steps decreased in October and January; however, not many changes were observed in the average exercise steps from 1200 to 1500 steps in all the months.

For the 6-months, the average walking steps were 4695.5±361.1, and the average exercise steps were 1394.8±183.4. In Klongyong, the average number of walking and exercise steps was 3741.3±129.1 and 1386.0±102.2. Overall, in Nihompattana, the walking steps (Klongyong: 3741.3±129.1 vs. Nihompattana; 4695.5±361.1,  $p < 0.001$ ) and exercise steps (Klongyong; 1386.0±102.2 vs. Nihompattana; 1394.8±183.4,  $p < 0.001$ ) were significantly higher than in Klongyong (Table 2).

### Anthropometry and blood pressure measurements

In Klongyong, a comparison of the anthropometry measurement results before and after the program showed that weight was not significantly different after the program. Notably, in Nihompattana, weight (before: 61.8 kg±1.1; after: 62.3 kg±1.1,  $p < 0.01$ ) and BMI (before: 24.9±0.4; after: 25.2 kg±0.4,  $p < 0.001$ ) significantly increased after the program. In Klongyong, a comparison of the blood pressure results from before and after the program showed that maximal blood pressure (before: 134.5 mmHg±4.1; and after: 126.0 mmHg±4.1,  $p < 0.05$ ) and minimal blood pressure (before: 77.2 mmHg±2.1; and after: 71.2 mmHg±2.3,  $p < 0.05$ ) showed significant decreases; however, in Nihompattana, no significant differences in maximal and minimal blood pressure were observed (Table 3).

### Physical fitness tests

In Klongyong, a comparison of the physical fitness test results from before and after the program showed no significant difference in sit-ups and eyes-open single-leg stance. The sit and reach flexibility (before:




Fig. 1: Daily steps and exercise steps, K: Klongyong, N: Nihompattana, ex: Exercise steps

Table 2: Comparison of the steps in Kloyong and Nihompattana

Walking steps	Kloyong	Nihompattana	p-value
Walking/day (steps)	3471.3±129.1	4965.5±361.1	0.000
Exercise/day (steps)	368.0±102.2	1394.8±183.4	0.006

49.9 cm±2.5; and after: 41.5±1.4,  $p < 0.001$ ) significantly decreased after the program. The hand grip strength (before: 19.9 kg±0.7; and after: 23.3 kg±1.0,  $p < 0.001$ ), 10-m obstacle walk (before: 13.0 s±1.2; and after: 9.4 s±0.6,  $p < 0.05$ ), and 6-min walk (before: 362.9 m±13.4; and after: 422.6 m±19.7,  $p < 0.01$ ) significantly improved after the program (Table 3).

In Nihompattana, a comparison of the physical fitness test results from before and after the program showed no significant difference in the number of sit-ups, 10-m obstacle walk, and 6-min walk. The hand grip strength (before: 26.3 kg±0.8; and after: 27.8 kg±0.7,  $p < 0.05$ ) and sit and reach flexibility (before: 34.4 cm±0.9; and after: 38.0 cm±1.3,  $p < 0.01$ ) significantly increased after the program. By contrast, eyes-open single-leg stance (before: 45.0 s±4.1; and after: 35.2 s±3.8,  $p < 0.01$ ) significantly decreased after the program. The sit and reach flexibility, 10 m-obstacle, and 6-min walk of interaction of the two-way ANOVA are significantly different (sit-and-reach flexibility;  $p < 0.001$ , 10 m-obstacle;  $p < 0.001$  and 6-min walk;  $p < 0.004$ ). Following the significant interaction, a one-way ANOVA was calculated to interpret the before and after results for the Klongyong sit and reach flexibility and showed a significant decrease (Klongyong before vs. Klongyong after;  $p < 0.05$ ); however, for the participants in Klongyong, the 10 m-obstacle, and 6-min walk showed significant improvement (Klongyong before vs. Klongyong after;  $p < 0.01$  and Klongyong before vs. Klongyong after;  $p < 0.05$ ) (Table 3).

### Brain function tests

In Klongyong, a comparison of the brain function in the go/no-go task from before and after the program showed no significant differences in formation reaction time, differentiation reaction time, reverse differentiation reaction time, and average reaction time. The total number of misses, mistakes, and errors was not significantly different after the program (Table 3).

In Nihompattana, a comparison of the brain function in the go/no-go task from before and after the program showed no significant differences in formation and average reaction times. Differentiation reaction times (before: 446.9 ms±9.0; and after: 481.7 ms±10.0,  $p < 0.001$ ) and reverse differentiation reaction times (before: 427.6 ms±12.1; and after: 446.2 ms±11.2,  $p < 0.001$ ) significantly increased after the program. By contrast, the total number of misses (before: 1.2 times±0.2; and after: 0.3 times±0.1,  $p < 0.001$ ), mistakes (before: 5.8 times±0.5; and after: 2.6 times±0.3,  $p < 0.001$ ), and errors (before: 7.1 times±0.6; and after: 2.9 times±0.3,  $p < 0.001$ ) significantly decreased after the program. The reverse differentiation reaction time, total number of misses, mistakes, and errors of interaction of the two-way ANOVA is significantly different

Table 3: Comparison of pre and post health program in Kloyong and Nihompattana

Measurement items	K/pre	K/post	p-value	N/pre	N/post	p-value	2 way ANOVA			Tukey-Kramer	
							B•A	K•N	Interaction	K/B•A	N/B•A
Body measurement											
Weight (kg)	55.0±2.3	54.7±2.5	0.482	61.8±1.1	62.3±1.1	0.006	0.743	0.001	0.870		
Body mass index	23.5±1.0	23.4±1.0	0.505	24.9±0.4	25.2±0.4	0.001	0.670	0.025	0.753		
Blood pressure											
Systolic blood pressure	134.5±4.1	126.0±4.1	0.042	134.0±1.7	135.7±1.6	0.271	0.981	0.093	0.082		
Diastolic blood pressure	77.2±2.1	71.2±2.3	0.022	79.2±1.3	79.0±1.0	0.889	0.327	0.010	0.160		
Physical fitness test											
Hand grip strength (kg)	19.9±0.7	23.3±1.0	0.001	26.3±0.8	27.8±0.7	0.002	0.036	0.001	0.372		
Sit up (times)	6.1±1.4	3.1±1.3	0.082	5.1±0.7	5.4±0.9	0.654	0.875	0.651	0.260		
Sit and reach flexibility (cm)	49.9±2.5	41.5±1.4	0.007	34.4±0.9	38.0±1.3	0.004	0.357	0.001	0.001	0.05	NS
Eyes open single leg stance (s)	13.9±2.6	20.4±5.6	0.167	45.0±4.1	35.2±3.8	0.002	0.168	0.001	0.167		
10 m obstacle walk (s)	13.0±1.2	9.4±0.6	0.030	8.5±0.2	8.4±0.2	0.079	0.054	0.001	0.001	0.01	NS
6 min walk (m)	362.9±13.4	442.6±19.7	0.002	514.3±9.6	508.3±8.7	0.778	0.336	0.001	0.004	0.05	NS
Brain function test (go/no-go task)											
Response											
Formation (ms)	585.9±56.3	745.5±56.3	0.178	356.4±13.5	380.0±14.4	0.119	0.111	0.001	0.092		
Differentiation (ms)	564.0±89.9	515.7±72.9	0.681	446.9±9.0	481.7±10.0	0.001	0.709	0.039	0.260		
Revers differentiation (ms)	566.0±71.3	493.3±40.8	0.391	427.6±12.1	446.2±11.2	0.001	0.495	0.011	0.030	NS	NS
Average (ms)	556.4±48.4	584.9±65.7	0.717	425.1±8.3	446.2±9.6	0.060	0.366	0.001	0.866		
Times											
Total number of misses (times)	0.0±0.0	0.0±0.0	-	1.2±0.2	0.3±0.1	0.001	0.001	0.001	0.016	NS	0.01
Total number of mistakes (times)	5.1±1.1	4.6±1.0	0.707	5.8±0.5	2.6±0.3	0.001	0.001	0.387	0.045	NS	0.01
Total number of errors (times)	5.1±1.1	4.6±1.0	0.707	7.1±0.6	2.9±0.3	0.001	0.001	0.859	0.002	NS	0.01

K: Kloyong, N: Nihompattana, B: Before, A: After, NS: Not significant

(reverse differentiation;  $p < 0.03$ , misses;  $p < 0.016$ , mistakes;  $p < 0.045$ , and errors;  $p < 0.002$ ). Following the significant interaction, a one-way ANOVA was calculated to interpret the before and after results for the reverse differentiation reaction time and showed no significant difference; however, in Nihompattana, the total number of misses, mistakes, and errors significantly decreased (Nihompattana before vs. Nihompattana after;  $p < 0.016$ , Nihompattana before vs. Nihompattana after;  $p < 0.045$ , and Nihompattana before vs. Nihompattana after;  $p < 0.002$ ) (Table 3).

## DISCUSSION

### Pedometry

Walking helps prevent diseases and contributes to health improvement [19,20]. In Klongyong, walking and exercise steps were approximately 3471.3 and 1386.0, respectively, and this result was lower than in Nihompattana, where the walking and exercise steps were approximately 4695.5 and 1394.8, respectively. An investigation in 2011 by the Japanese Ministry of Health, Labor, and Welfare showed that the average number of steps was 7841 for men and 6883 for women aged 20–64 years, and aims for 9000 steps for men and 8500 steps for women by 2020 [21].

### Anthropometry and blood pressure measurements

In Nihompattana, a comparison of the anthropometry results from before and after the program showed that weight significantly increased after the program. In Klongyong, weight increased after the program. In addition, in Nihompattana, maximal and minimal blood pressures from before and after the program showed no significant differences. By contrast, in Klongyong, maximal and minimal blood pressures showed significant decreases. Regarding the health education provided in Japan, participants' weight, BMI, and blood pressure tended to decrease [12,13]. The significant increase in weight and BMI of the participants' in Nihompattana maybe because of the shorter duration of the health education enforcement period.

### Physical fitness tests

In Klongyong, a comparison of the physical fitness test results from before and after the program showed significant improvements in the hand grip strength, 10 m obstacle walk, and 6-min walk; however, sit and reach flexibility significantly decreased after the program. In Nihompattana, the hand grip strength and sit and reach flexibility significantly improved after the program; however, eyes-open single-leg stance significantly decreased after the program. The difference

in the results may be the effects of the age difference; that is, the Klonyong participants are approximately 10 years older than those in Nihompattana.

In Klonyong, the physical fitness test results showed that sit-and-reach flexibility significantly decreased. In Nihompattana, eyes-open single-leg stance significantly decreased after the program. Increased exercise momentum leads improved physical fitness test results [22].

For Klonyong and Nihompattana, it is desirable to exercise with the goal of walking 7000 steps [11-14]. The 10 m-obstacle walk and 6-min walk of the interaction of the two-way ANOVA suggest that the participants in Klonyong showed significant improvement. It is possible that the significantly increased rate of improvement in Klonyong is because its initial values were higher than in Nihompattana.

### Brain function tests

Go/no-go tasks are frequently used to investigate response inhibition, an essential executive function implemented by the prefrontal cortex, and these tasks recruit a variety of cognitive components in addition to response inhibition [16-18,23,24]. In Nihompattana, the total number of misses, mistakes, and errors significantly improved after the program. Go/no-go task studies in the literature have suggested that a health program could improve brain function, including working memory [24-26]; in those studies, participants performed regular exercises in the first stage, where go/no-go task reaction times increased and number of error responses decreased significantly. In the second stage, go/no-go task reaction times decreased significantly, and the number of error responses decreased significantly [11-14].

The results from Klonyong suggest that, in the first stage, the go/no-go task reaction time increased and number of error responses decreased. Notably, the results from Nihompattana suggest that, in the first stage, the go/no-go task reaction time increased and number of error responses significantly improved. In other words, the total number of misses, mistakes, and errors of the interaction of the two-way ANOVA suggests that the Nihompattana showed significant improvement. These results may be due to the greater amount of exercise performed by the Nihompattana participants. The number of daily mean steps was 1200 more than in Klonyong. It has been suggested that a walk and a walk with dual-task improve brain function [25,26]. The quantity of the exercise might induce the improvement of the brain function [27]. Notably, the initial values were higher in Klonyong than in Nihompattana. This result may be the reason for the significantly increased rate of improvement in Klonyong.

### The limitations of this study

The Japanese Ministry of Education, Culture, Sports, Science, and Technology recommends that Japanese individuals, from elementary school students to elderly people, should measure their physical fitness. Therefore, they are used to taking physical fitness tests. However, elderly people in Thailand have never experienced doing so and, therefore, may not easily understand how to do so. This factor may be one of the causes of Thailand's lower values in the test results. In the future, it is necessary to increase the number of subjects, to implement health education in more areas, and to find the improvement points of the health education system.

### CONCLUSIONS

We implemented a 12-month health program in Klonyong and a 6-month health program in Nihompattana. This health promotion program used the international organization for standardization assessed findings from pedometer, anthropometry, blood pressure, physical fitness, and brain function. In Klonyong, walking and exercise steps were approximately 3471.3 and 1386.0, respectively, which was less than in Nihompattana, where the walking and exercise steps were approximately 4695.5 and 1394.8, respectively.

In Klonyong, a comparison of the physical fitness test results from before and after the program showed significant improvements in the hand grip strength, 10 m obstacle walk, and 6-min walk; however, sit and reach flexibility significantly decreased after the program. In Nihompattana, hand grip strength and sit and reach flexibility significantly improved after the program; however, eyes-open single-leg stance significantly decreased after the program. The 10m-obstacle walk and 6-min walk of the interaction of the two-way ANOVA suggest that the Klonyong showed significant improvement, compared with Nihompattana; however, the significantly increased rate of improvement is because the initial values of Klonyong were higher than those of Nihompattana. The total number of misses, mistakes, and errors of the interaction of the two-way ANOVA suggests that the Nihompattana showed significant improvement, compared with Klonyong. These results may be due to the greater amount of exercise performed by the Nihompattana participants. The number of the daily mean steps was 1200 more than in Klonyong. This result suggested that a walk and a walk with dual-task improve brain function [23,24]. The quantity of the exercise might induce the improvement in brain function [25]. Notably, the initial values of the participants in Klonyong were higher than in Nihompattana; thus, this phenomenon may have led to the significantly increased rate of improvement in Klonyong.

### AUTHOR'S CONTRIBUTIONS

SJ, KT, NP, SK, PU, SS, TW, and NT planned the experiment and collected the data. MO, HT, HK, RU, and KA conducted data analysis. SJ, KT, and HT formulated the composition of the article. All authors read and approved the final manuscript.

### AUTHORS FUNDING

The authors thank the individuals who participated in this study. Suchinda Maruo Jarupat is supported by Thai Population Development Project of the Faculty of Medicine, Ramathibodi Hospital, Mahidol University, Thailand. Koji Terasawa is supported by a Grant-in-Aid for the Scientist (Houga:26560380, Kiban A:25257101) of the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

### CONFLICTS OF INTEREST

There are no conflicts of interest.

### REFERENCES

1. Declaration of ALMA-ATA. *Am J Public Health* 2015;105:1094-5.
2. Ilona K. Health Promotion a Discussion Document on the Concept and Principles of Health Promotion. World Health Organization Regional Office for Europe; 1984. p. 73-6.
3. Adepu SP, Sam S, Omanakuttan C, Ramanatha KV, Yashaswini Y. Assessment of pharmacist mediated education on medication adherence behavior in Type 2 diabetes mellitus patients in South Indian rural population. *Int J Pharm Pharm Sci* 2018;10:39-43.
4. Missriya MS, Subashini S. Assess the prevalence of respiratory problem and knowledge on health effects of people residing near quarry. *Int J Pharm Pharm Sci* 2017;9:250-3.
5. Kawamori M. Community Health-oriented Bureaucrats and the Development of Thai Health Policy Individuals, Organization and Movements in Health System Reform. Geneva: World Health Organization; 2006. p. 1-19.
6. World Economics Forum. Human Capital Outlook: Association of Southeast Asian Nations (ASEAN). Regional Country Briefing; 2016. p. 1-12.
7. Greider AC, Knebel E. Health Professions Education. United States: National Academies Press; 2003. p. 29-67.
8. Nola P, Carolyn M, Mary P. Health Promotion in Nursing Practice. New Jersey: Person Prentice Hall; 2006. p. 74-91.
9. Piyanun L, Nittaya S, Ameporn R. Effectiveness of a self-regulation program on diet control, exercise, and two-hour postprandial blood glucose levels in Thais with gestational diabetes mellitus. *Pac Rim Int J Nurs Res* 2011;15:173-86.
10. Nongyaow M, Linchong P, Sharon K, Ostwald S. Effectiveness of nutritional education in promoting healthy eating among elders in Northeastern Thailand. *Pac Rim Int J Nurs Res* 2011;15:88-201.

11. Nakade K, Abe K, Fujiwara T, Terasawa K, Okuhara M, Nakajima K, *et al.* The influence of two different health education program on GO/NO-GO tasks, physical fitness tests and blood tests. *Jpn Soc Phys Anthropol* 2009;14:143-50.
12. Murata Y, Nemoto K, Kobayashi I, Miyata Y, Terasawa S, Sasamori F, *et al.* Effect of a two-year health program on brain function, physical fitness and blood chemistry. *J Community Med Health Educ* 2015;5:1-6.
13. Watanabe T, Terasawa K, Nakade K, Murata Y, Terasawa S, Nome K, *et al.* Differences between two Japanese health-promotion programs on measures of health and wellness. *J Med Health Sci* 2015;5:170-81.
14. Suchinda JM, Sanongdetch W, Choeychom S, Chansirikarn S, Thrakul S, Phuphaibul R, *et al.* Comparing the effectiveness of health program in Thailand and Japan. *J Nurs Care* 2015;4:1-6.
15. Ministry of Education, Culture, Sports, Science and Technology; 2019. Available from: [https://www.mext.go.jp/component/a\\_menu/sports/detail/\\_icsFiles/afieldfile/2010/07/30/1295079\\_04.pdf](https://www.mext.go.jp/component/a_menu/sports/detail/_icsFiles/afieldfile/2010/07/30/1295079_04.pdf). [Last accessed on 2019 Dec 20].
16. Masaki T, Moriyama G. Study on Types of Human Higher Nervous Activity. Vol. 4. Tokyo: University of Science Press; 1971. p. 69-81.
17. Terasawa K, Tabuchi H, Yanagisawa H, Yanagisawa A, Shinohara K, Terasawa S, *et al.* Comparative survey of Go/no-go results to identify the inhibitory control ability change of Japanese children. *Biopsychosoc Med* 2014;14:1-8.
18. Terasawa K, Misaki S, Murata Y, Watanabe T, Terasawa S, Kobayashi T, *et al.* Relevance between Alzheimer's disease patients and normal subjects using Go/no-go tasks and Alzheimer assessment scores. *J Child Adolesc Behav* 2014;2:1-5.
19. Hamer M, Chid Y. Walking and primary prevention: A meta-analysis of prospective cohort studies. *Br J Sports Med* 2008;42:238-43.
20. Hornestam JF, Souza TR, Arantes P, Ocarino J, Silva PL. The Effect of walking speed on foot kinematics is modified when increased pronation is induced. *J Am Podiatr Med Assoc* 2016;106:419-26.
21. Ministry of Health, Labor and Welfare; 2019. Available from: <https://www.mhlw.go.jp/topics/2009/05/dl/tp0501-siryous3-5.pdf>. [Last accessed on 2019 Dec 20].
22. Leite JC, Forte R, Vito DG, Boreham CA, Gibney MJ, Brennan L, Gibney ER. Comparison of the effect of multicomponent and resistance training programs on metabolic health parameters in the elderly. *Arch Gerontol Geriatr* 2015;60:412-7.
23. Diamond A. Executive functions. *Annu Rev Psychol* 2013;64:135-68.
24. Chikazoe J. Localizing performance of go/no-go tasks to prefrontal cortical subregions. *Curr Opin Psychiatry* 2010;23:267-72.
25. Toots A, Littbrand H, Holmberg H, Nordström P, Lundin-L LL, Gustafson Y, *et al.* Walking aids moderate exercise effects on gait speed in people with dementia: A randomized controlled trial. *J Am Med Dir Assoc* 2016;16:30402-9.
26. Nakade K, Watanabe T, Miura H, Murata Y, Terasawa S, Kobayashi T, *et al.* Effect of Training on Dementia Prevention while Performing a Dual Task during Exercise. Bangkok, Thailand: Presented at the 6<sup>th</sup> International Society for Physical Activity and Health; 2017.
27. Babaei P, Alamdari AK, Soltani TB, Damirchi A. Effect of six weeks of endurance exercise and following detraining on serum brain derived neurotrophic factor and memory performance in middle aged males with metabolic syndrome. *J Sports Med Phys Fitness* 2013;53:437-43.