

NATURAL FLAVONOIDS: A NOVEL APPROACH TO BREAST CANCER (REVIEW)

SHIKHA RANA¹, SAVITA DIXIT*, ALOK MITTAL**

1. Research scholar, Maulana Azad Institute of Technology, Bhopal. *Professor, HOD, Department of Chemistry, Maulana Azad Institute of Technology, Bhopal. ** Professor, Department of Chemistry, Maulana Azad Institute of Technology, Bhopal.
Email: Savitadixit1@yahoo.com

Received: 10 March 2017, Revised and Accepted: 20 May 2017

ABSTRACT

Cancer is a hereditary disease and is caused due to the abnormal growth of the cells. Cancer can be of many forms but the most prevalent is the breast cancer. Breast cancer is the most common among women and can be treated by radiation therapy, surgery, hormone therapy, chemotherapy and natural therapy. The present review presents an overview about the role of flavonoids in curing cancer. This review gives a detailed account of classification of flavonoids. The in vivo and in vitro studies show the action of flavonoids on cancer. Various natural flavonoids can be extracted and can be used for various therapeutic effects apart from cancer.

Keywords: Flavanoids, Cancer, Breast Cancer, Classification of flavanoids.

INTRODUCTION

Cancer is a hereditary disease. The agents causing cancer are known as carcinogens. These can be present in food, water and air. It is a virulent disease which leads to the genetic mutations bringing a change in regulating proteins. The resultant abnormal cell behaviour leads to extensive growth of cells affecting various surrounding cells and organs resulting into cancer and ultimately leads to the death of the patient (1).

Breast Cancer is the most prevailing disease in women. It is the second common disease after lung cancer. The disclosure, medication and avoidance of breast cancer are one of the burning issues in public health and medical practices (2).


Breast Cancer patients can be diagnosed by surgery, radiation therapy, chemotherapy, hormone therapy and natural therapy. These therapies are employed to wipe out cancer, avert the production of metastases and counter the exoneration (3).

Historical Perspective

A new flavonoid called rutin was isolated from oranges in the year 1930 which reduced the capillary permeability (35). Flavonoids are found in fruits, vegetables, grains, bark, roots, stems, flowers, tea and wine (36). Flavonoids are found in various medicinal plants and find a great use in various treatments used worldwide specially in China. Flavonoids extracted from licorice have been found to show their therapeutic effects against peptic ulcers and gastric cancer in *H. pylori*-infected individuals (37). Genistein present in soy was tested with various other natural flavonoids and was found to inhibit cell proliferation in estrogen receptor positive breast cancer cells. Another such flavonoid is Quercetin which is found effective against breast cancer and is effective only in water soluble form (38). It was also found in cell culture studies that many flavonoids could inhibit breast cancer resistance protein (BCRP, ABCG2) (39). Flavonoids are also reported to show chemopreventive effects in estrogen dependent or independent breast cancer (40). Many fruits and leafy vegetables are found to contain kaempferol. Onion and pears are found to contain isorhamnetin whereas myricetin is found in berries, maize and tea. Citrus fruits and vegetables are rich sources of anthocyanidins. Soya and soya products mainly include daidzein and genistein which are the isoflavonones (41). Black tea is the rich source of flavan-3-ols. The red skin peanuts are also reported to contain some amount of flavonones.

Overview (Flavonoids)

Flavonoids are the subclass of polyphenols which are extensively dispersed in plants such as citrus fruits, berries, onions, parsley, legumes, green tea, red wine, sea buckthorn and dark chocolate. It consists of a diphenylpropane skeleton, consisting of two aromatic rings (i.e. A-ring and B-ring), each contains at least one hydroxyl group, and the two aromatic rings are connected through a three-carbon bridge, which becomes part of the six-member heterocyclic ring (4).




On the basis of their chemical structures flavonoids are classified into following (Table 1):

1. flavanoids (2-phenylbenzopyrans)
2. isoflavonoids (3-benzopyrans)
3. neoflavonoids (4-benzopyrans) (5,6)


2.1 Flavonoids

These are further divided into the following depending upon degree of oxidation and saturation present in heterocyclic ring.



2.2 Isoflavonoids

These are also of following categories:



2.3 Neoflavonoids

These comprises of the following:




Table 1.


S No	Groups	Sub Groups
1.	Flavonoids	Flavan
		Flavanone
		Flavone
		Flavonol
		Dihydroflavonol
		Flavon-3-ol
		Flavon-4-ol
Flavan-3,4-diol		
2.	Isoflavonoids	Isoflavan
		Isoflavone
		Isoflavanone
		Isoflav 3 ene
		Isoflavanol
		Rotenoid
		Coumestane
		3 aryl coumarin
		Coumaronochremene
		Coumaronochromone
Pterocarpan		
3.	Neoflavonoids	4- aryl coumarin
		3,4 - dihydro-4- aryl coumarin
		Neoflavene

3,Flavonoids and Cancer

Flavonoids display an exceptional spectrum of biological activities which may affect cancer activities. These can be antiallergic, anti-inflammatory, antioxidant, antimutagenic, anticarcinogenic, and modulation of enzymatic activities (7-9).These are benign in cancer chemoprevention and may acts as potential therapeutic agents (10).

Carcinogenesis is a multistep process. Its steps can be summarised as

- Initiation: It is an accelerated phase which includes interaction of DNA with carcinogenic material.
- Promotion: The abnormal cells flourish and grow.
- Progression:It is the final stage in which premalignant cells into neoplastic cells (11).



Flavonoids are most widely found in photosynthesising plant cells. These are the indispensable part in human and animal diet and cannot be synthesised within body. In recent years there has been an upswing concern in the therapeutic potential of flavonoids. These are mainly due to the presence of phenolic groups (12, 13). Various edible plants have been linked to treatment of cancer (14). Various plant derived agents such as paclitaxel, docetaxel; vinblastine, vincristine; topotecan, irinotecan, etoposide are currently being used for treatment of cancer (15, 16, 17).

Plants have various flavonoids which are biologically very active and possess various therapeutic properties which outlines its necessity for determination (18). Flavonols such as catechin, epicatechin and epigallocatechin from tea have been extracted(19). Certain flavones such as Chrysin, apigenin, Rutin, luteolin, and luteolin glucosides are found in the fruit skins, red wine, buckwheat, red pepper, and tomato skin(20-23). Flavonols such as Kaempferol, quercetin, myricetin, and tamarixetin are found in Onion, red wine, olive oil, berries, and grapefruit (22). Citrus fruits, grapefruits, lemons, and oranges possess flavonone such as naringin, naringenin, taxifolin, and hesperidin(24, 25). Soybean consists of isoflavone such as genistin, daidzin(26).

Polyphenols found in tea especially in green tea have shown to reduce the risk of cancer (27). Curcumin in turmeric has been found to affect various multicellular signalling pathways which are involved in proliferation, invasion, survival, apoptosis and inflammation (28). Various other plants and their phytochemicals effective against cancer are listed in the following table.

SNo	Source	Phytochemicals	Reference
1.	broccoli, sprouts, cabbage, kale	Sulforaphane	29
2.	broccoli, cabbage, cauliflower, mustard, and radish	Indole-3 carbinol, diindolylmethane	30
3.	Soy Compounds	Genistein, Soy Isoflavones	31
4.	wasabi, horseradish, mustard, radish, brussel sprouts, watercress, nasturtiums, capers	isothiocyanate	32
5.	red wine, peanuts, and certain berries	Resveratrol	33
6.	garlic, chives, and leeks	Organosulphur Compounds	27
7.	tomato products.	Lycopene	35
8.	citrus fruits and buckwheat	Quercetin	

3.1 In vitro studies

Many researchers have conducted various in vitro studies on the potential anticancer activity of flavonoids. Biphase effects of isoflavones have been seen in proliferation of breast cancer cell culture. At concentration above 5mM genistein showed concentration dependant ability to inhibit growth and estrogen stimulated breast cancer cell proliferation. 28 flavonoids were studied by Hirano and co workers against acute myeloid cell line HL 60 and was compared with antiproliferative activity and cytotoxicity with four clinical anticancer agents. Out of these 28 flavonoids eight shows suppressive effects on HL 60 cell growth. The rest had potent anticancer activity (42). 55 flavonoids were evaluated by Cushman and Nagarthnam and studied their toxicity in five cancer cell cultures and were found effective against A-549 lung carcinoma, MCF - 7 breast carcinoma, HT - 29 colon adenocarcinoma, SKMEL- 5 melanoma and MLM melanoma (43). 27 out of these were of citrus origin and were found to inhibit tumor cell proliferation.

3.2 In vivo studies

Flavonoids have also shown to exhibit anticancerous activities in vivo studies. They may inhibit carcinogenesis in any stage of carcinoma. Animal and other cellular model investigations showed that certain flavonoids inhibit tumor initiation and progression. A novel research showed that fermented soy milk contains large amounts of genistein and daidzein which were given to rats of 7 weeks of age showed to inhibit mammary tumorigenesis which was induced artificially by 2-amino-1-methyl-6-phenylimidazo [4,5-b] pyridine (PhIP) (44). Mammographic breast density can be used as biomarkers for estrogenic and anti estrogenic effects of a particular treatment in breast tissue. Consuming dietary supplements including red clover derived isoflavones for 12 months did not show any increase in mamographic breast density in women experiencing menopause suggesting that there is no effects of estrogenic and anti estrogenic effects on breast (45).

CONCLUSION

Flavonoids have a wide range of therapeutic effects. The impact of flavonoids has been seen widely during the avalanche of immunological events which are correlated with the advancement and progression of cancer. A methodical mechanism is to be developed so as to understand mechanism of action of flavonoids when they enter various cell organelles and tissues. Flavonoids are probable to inflect various biological events in cancer including apoptosis, vascularization, cell differentiation, cell proliferation etc. Various dietary flavonoids show anti tumor activity during in vivo studies whereas these shows repression in vivo studies. Various potent flavonoids are to be studied and extracted for elucidating various other natural ways in treating cancer. Further studies should be conducted so as to validate the traditional ways of treating cancer. In the past many efforts are made to get various anticancerous plants containing flavonoids and further studies are to be made to get satisfactory results. For these a number of medicinal plants can be screened and can be further worked on so that in vitro and in vivo studies can be conducted providing new insights for fighting against cancer.

REFERENCES

1. Cancer; Malcolm R Alison, Imperial College School of Medicine, London, UK. ENCYCLOPEDIA OF LIFE SCIENCES / & 2001 Nature Publishing Group / www.els.net.
2. Breast Cancer and the Environment A Life Course Approach, http://iom.nationalacademies.org/~media/Files/Report%20Files/2011/Breast-Cancer-Environment/breastcancerreportbrief_2.pdf
3. Breast Cancer Treatment Options: Surgery and Radiation Therapy: A Literature Review. Breast cancer treatment options: a literature review, December 2012
4. Effects of Flavonoids on Proliferation of Breast Cancer Cells and Vascular Smooth Muscle Cells Liu Po Shiu Jackie, A thesis submitted for the degree of Master of Medical Sciences (2007), THE UNIVERSITY OF HONG KONG.
5. [Http://files.rushim.ru/books/mechanizms/the-science-of-flavonoids.pdf](http://files.rushim.ru/books/mechanizms/the-science-of-flavonoids.pdf)
6. The Science of Flavonoids, THE STEREOCHEMISTRY OF FLAVONOIDS, Jannie P.J. Marais, Bettina Deavours, Richard A. Dixon, and Daneel Ferreira. Springer, ISBN-13: 978-0387-28821-5, 2006 Springer Science Business Media, Inc.
7. Middleton EJ, Kandaswami C, Theoharides TC. The effects of plant flavonoids on mammalian cells: Implications for inflammation, heart disease and cancer. *Pharmacol Rev.* 2000;52:673-751. Pubmed
8. Galati G, Teng S, Moridani MY, Chan TS, O'Brien PJ. Cancer chemoprevention and apoptosis mechanisms induced by dietary polyphenolics. *Drug Metabol Drug Interact.* 2000;17:311-49. Pubmed
9. Yang CS, Landau JM, Huang MT, Newmark HL. Inhibition of carcinogenesis by dietary polyphenolic compounds. *Annu Rev Nutr.* 2001;21:381-406. Pubmed
10. Birt DF, Hendrich S, Wang W. Dietary agents in cancer prevention Flavonoids and isoflavonoids. *Pharmacol Ther.* 2001; 90:157-77. Pubmed
11. Flavonoids: A versatile source of anticancer drugs, Maheep K Chahar, Neelu Sharma, Mahabeer P Dobhal, Yogesh C Joshi, Review Article, Year: 2011, Volume: 5, Issue: 9, Page: 1-12. DOI: 10.4103/0973-7847.79093.PMC
12. Pourmorad F, Hosseinimehr SJ, Shahabimajd N. Antioxidant activity, phenol and flavonoid contents of some selected Iranian medicinal plants. *The African Journal of Biotechnology.* 2006;5(11):1142-1145.
13. Kumar S, Pandey AK. Antioxidant, lipo-protective and antibacterial activities of phytoconstituents present in *Solanum xanthocarpum* root. *International Review of Biophysical Chemistry.* 2012;3(3):42-47.
14. Mignet N, Seguin J, Ramos Romano M, Brulle L, Touil YS, Scherman D, et al. Development of a liposomal

- formulation of the natural flavonoid fisetin. *Int J Pharm.*2012;423:69–76. Pubmed
15. Kim JH, Jung CH, Jang BH, Go HY, Park JH, Choi YK, et al. Selective cytotoxic effects on human cancer cell lines of phenolic-rich ethyl-acetate fraction from *Rhus verniciflua* Stokes. *Am J Chin Med.* 2009;37:609–20. Pubmed
 16. Li YL, Gan GP, Zhang HZ, Wu HZ, Li CL, Huang YP, et al. A flavonoid glycoside isolated from *Smilax china* L. Rhizome *in vitro* anticancer effects on human cancer cell lines. *J Ethnopharmacol.* 2007;113:115–24. Pubmed
 17. Attoub S, Hassan AH, Vanhoecke B, Iratni R, Takahashi T, Gaben AM, et al. Inhibition of cell survival, invasion, tumor growth and histone deacetylase activity by the dietary flavonoid luteolin in human epithelioid cancer cells. *Eur J Pharmacol.* 2011;651:18–25. Pubmed
 18. Total Phenolics and Total Flavonoids in Selected Indian Medicinal Plants, C. T. Sulaiman* and Indira Balachandran, *Indian J Pharm Sci.* 2012 May-Jun; 74(3): 258–260.PMC.
 19. Lopez M, Martinez F, Del Valle C, Orte C, Miro M. Analysis of phenolic constituents of biological interest in red wines by high-performance liquid chromatography. *Journal of Chromatography A.*2001;922(1-2):359–363. [pubmed]
 20. Hara Y, Luo SJ, Wickremasinghe RL, Yamanishi T. Special issue on tea. *Food Reviews International.*1995;11:371–542.
 21. Kreft S, Knapp M, Kreft I. Extraction of rutin from buckwheat (*Fagopyrum esculentum* Moench) seeds and determination by capillary electrophoresis. *Journal of Agricultural and Food Chemistry.*1999;47(11):4649–4652. [pubmed]
 22. Stewart AJ, Bozonnet S, Mullen W, Jenkins GI, Lean ME, Crozier A. Occurrence of flavonols in tomatoes and tomato-based products. *Journal of Agricultural and Food Chemistry.* 2000;48(7):2663–2669.[pubmed]
 23. Hertog MGL, Hollman PCH, Katan MB. Content of potentially anticarcinogenic flavonoids of 28 vegetables and 9 fruits commonly consumed in the Netherlands. *Journal of Agricultural and Food Chemistry.* 1992;40(12):2379–2383.
 24. Miyake Y, Shimoi K, Kumazawa S, Yamamoto K, Kinase N, Osawa T. Identification and antioxidant activity of flavonoid metabolites in plasma and urine of eriocitrin-treated rats. *Journal of Agricultural and Food Chemistry.* 2000;48(8):3217–3224. [pubmed]
 25. Rousseff RL, Martin SF, Youtsey CO. Quantitative survey of narirutin, naringin, hesperidin, and neohesperidin in citrus. *Journal of Agricultural and Food Chemistry.* 1987;35(6):1027–1030.
 26. Reinli K, Block G. Phytoestrogen content of foods: a compendium of literature values. *Nutrition and Cancer.* 1996;26(2):123–148. [pubmed]
 27. Plant Phytochemicals as Epigenetic Modulators: Role in Cancer Chemoprevention, Vijay S. Thakur, Gauri Deb, Melissa A. Babcock, and Sanjay Gupta. *AAPS J.* 2014 Jan; 16(1): 151–163.PMC
 28. Kunnumakkara AB, Anand P, Aggarwal BB. Curcumin inhibits proliferation, invasion, angiogenesis and metastasis of different cancers through interaction with multiple cell signaling proteins. *Cancer Lett.*2008;269:199–225. [pubmed]
 29. Multi-targeted prevention of cancer by sulforaphane (review). Clarke JD, Dashwood RH, Ho E *Cancer Lett.* 2008 Oct 8; 269(2):291-304.[pubmed]
 30. Banerjee S, Kong D, Wang Z, Bao B, Hillman GG, Sarkar FH. Attenuation of multi-targeted proliferation-linked signaling by 3,3'-diindolylmethane (DIM): from bench to clinic. *Mutat Res.*2011;728:47–66. [PMC free article] [pubmed]
 31. Multi-targeted therapy of cancer by genistein (review). Banerjee S, Li Y, Wang Z, Sarkar FH *Cancer Lett.* 2008 Oct 8; 269(2):226-42. [pubmed]
 32. Review, Molecular targets of dietary phenethyl isothiocyanate and sulforaphane for cancer chemoprevention. Cheung KL, Kong AN *AAPS J.* 2010 Mar; 12(1):87-97(PUBMED)
 33. Resveratrol and cancer: a review. Savouret JF, Quesne M *Biomed Pharmacother.* 2002 Mar; 56(2):84-7. (PMC)
 34. Tomatoes, tomato-based products, lycopene, and cancer: review of the epidemiologic literature. Giovannucci E *J Natl Cancer Inst.* 1999 Feb 17; 91(4):317-31.(Pubmed)
 35. Wine, alcohol, platelets, and the French paradox for coronary heart disease. Renaud S, de Lorgeril M *Lancet.* 1992 Jun 20; 339(8808):1523-6 pubmed.
 36. Flavonoids: a review of probable mechanisms of action and potential applications. Nijveldt RJ, van Nood E, van Hoorn DE, Boelens PG, van Norren K, van Leeuwen PA *Am J Clin Nutr.* 2001 Oct; 74(4):418-25. (pubmed)
 37. Anti-helicobacter pylori flavonoids from licorice extract. Fukai T, Marumo A, Kaitou K, Kanda T, Terada S, Nomura T. *Life Sci* 2002; 71:1449–1463.
 38. [Http://www.allcancerguide.com/2015/07/flavonoids-in-breast-cancer.html](http://www.allcancerguide.com/2015/07/flavonoids-in-breast-cancer.html)
 39. [Http://dmd.aspetjournals.org/content/33/3/341.full](http://dmd.aspetjournals.org/content/33/3/341.full)
 40. Breast Cancer and Flavonoids - A Role in Prevention. Hitomi Takemura, Hiroyuki Sakakibara, Shunsuke Yamazaki and Kayoko Shimoi *Current Pharmaceutical Design*, 19(34):6125-6132.
 41. Flavonoid intake and breast cancer risk: a case-control study in Greece. J Peterson et al. *Br J Cancer.* 2003 Oct 6; 89(7): 1255–1259.
 42. Natural flavonoids and lignans are potent cytostatic agents against human leukemic HL-60 cells. Hirano T, Gotoh M, Oka K. *Life Sci* 1994;55:1061–1069.
 43. Cytotoxicities of some flavonoid analogues. Cushman M, Nagarathnam D. *J Nat Prod* 1991;54:1656–1660.
 44. Inhibitory effects of bifidobacterium-fermented soy milk on 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine-induced rat mammary carcinogenesis, with a partial contribution of its component isoflavones. Ohta T, Nakatsugi S, Watanabe K, Kawamori T, Ishikawa F, Morotomi M, Sugie S, Toda T, Sugimura T, Wakabayashi K *Carcinogenesis* 2000;21:937–941.
 45. Anti-cancer potential of flavonoids: recent trends and future perspectives. Priya Batra• Anil K. Sharma. *Review, 3 Biotech(2013), 3:439-459.*