

## PHARMACOGNOSTICAL, PHYTOCHEMICAL STUDIES ON THE LEAVES OF *Psidium guajava* Linn- LUCKNOW- 46 VARIETY

**K.Periyanayagam\*, S.Gopalakrishnan, V.Karthikeyan**

Asst Reader, Department of Pharmacognosy, College of pharmacy, Madurai Medical College, Madurai 625 020, Tamil nadu, India.

Email: kpn1960@yahoo.com

Received: 6 September 2013, Revised and Accepted: 30 September 2013

### ABSTRACT

**Objective:** To explore the micro morphology and physio chemical parameters of the leaves of *Psidium guajava* Linn. (Myrtaceae) – Lucknow- 46 variety.

**Methods:** Macroscopy, microscopy, physicochemical analysis, preliminary phytochemical screening and other WHO recommended parameters for standardizations were performed.

**Results:** Leaves (5-15cm × 4-6 cm) are dorsiventral, oblong – elliptic, dull grey to yellow green with entire margin, obtuse to bluntly acuminate apex and rounded to subcuneate base with short petiole. Microscopic evaluation revealed the presence of paracytic stomata, three layered thin walled tabular epidermal cells and wider spindle shaped sub epidermal layers with frequent calcium oxalate druses, three layers of narrow palisade cells, vertically stretched collateral vascular bundles with sclerenchyma bundle caps, spongy parenchyma is not distinct but contains abundant tannin content, circular secretory cavities, calcium oxalate druses in ground tissue, broadly bowl shaped vascular bundles, xylem vessels, phloem, and fibres. Vein islet numbers, vein termination numbers, stomatal number, stomatal index and other physico chemical tests like ash values, loss on drying, extractive values were determined. Preliminary phytochemical screening showed the presence of sterols, tannins, proteins and aminoacids, flavonoids, volatile oil, terpenoids, saponin, carbohydrates and absence of alkaloids, mucilage, glycosides, fixed oil.

**Conclusion:** Microscopic analysis was informative and provides useful information in the botanical identification, standardization for purity & quality and immense value in authentication of the leaf.

**Keywords:** *Psidium guajava*, Myrtaceae, Microscopical evaluation, Physicochemical analysis.

### INTRODUCTION

*Psidium guajava* Linn commonly called as poor man apple. The leaves of *P. guajava* really do not have any match as a cheap natural and easily available plant. It is traditionally known to be useful for the treatment of wide panel of diseases like ulcers, wounds, astringent, antiemetic, cholera, epilepsy etc [1]. Leaf is traditionally used for antispasmodic, anodyne, febrifuge[2], scurvy [3], malaria [4], antiseptic[5], antibacterial[6-8], antifungal [9] dysentery, diarrhoea[10,11], anti-inflammatory [12,13], gout[14], hypoglycaemic[15], headache, fever, gonorrhoea, dysmenorrhoea[16], haemostat[17], antihypertensive[18], analgesic[19], hepatoprotective[20] and anticoagulant[21].

It was reported that fresh leaves contains: Guajavarin, isoquercetin, hyperin, quercetrin, quercetin 3-o gentiobioside [22]. Leaves also contains two triterpenoids, guavanoic acid and guava coumaric acid along with six known compounds 2 alpha hydroxy ursolic acid, jacoumaric acid, isoneriucoumaric acid, asiatic acid, ilelatifol D and β-sitosterol – 3-o – beta D glucopyranoside [23]. In short, there is good level of traditional and experimental evidences to support various claims and advantages of this widely available plant. An investigation to explore its pharmacognostic examination is inevitable. Hence, in this work we report an attempt on microscopic evaluation, physicochemical determination and phytochemical screening for the standardization and quality assurance purposes of this cultivar.

### MATERIALS AND METHODS

#### Chemicals

Formalin, acetic acid, ethyl alcohol, chloral hydrate, toluidine blue, phloroglucinol, glycerin, hydrochloric acid and all other chemicals used in this study were of analytical grade.

#### Plant collection and authentication

The leaves of the healthy plant *Psidium guajava* Linn. (Lucknow – 46) selected for our study was collected from Horticulture Department, Madurai, Tamilnadu, India and was authenticated by **Dr. Stephen**, Department of Botany, American college, Madurai and **Dr. P. Jayaraman**, Director of Plant Anatomy Research Institute, Tambaram, Chennai, Tamil nadu, India.

#### Macroscopic analysis

Macroscopic observation of the plant was done. The shape, size, surface characters, texture, colour, odour, taste etc was noted [24].

#### Microscopic analysis

Transverse section midrib region of fresh leaf pieces were cut and fixed in FAA and then dehydrated by employing graded series of ethyl alcohol and tertiary butyl alcohol [25]. Sections were taken using microtome. Permanent mount was prepared using safranin fast green double staining technique [26]. In order to supplement the descriptive part the photomicrographs in different magnifications of all necessary cells and tissues were taken with NIKON Coolpix 8400 digital camera and Labphot2 microscopic unit.

#### Powder microscopy

Coarse powder of the leaf was used to study the microscopical characters of the leaf powder [27,28].

#### Physicochemical analysis

Total ash, acid insoluble ash, water soluble ash, loss on drying, extractive values and leaf constants such as vein islet numbers, vein

terminal number, stomatal number and stomatal index, palisade ratio were determined [29-31].

#### Preliminary phytochemical screening:

Preliminary phytochemical screening was carried out to find out the presence of various phytoconstituents using standard procedure [32, 33].

### RESULTS

#### Macroscopy

*Psidium guajava* is a large dicotyledonous- shrub or small evergreen tree, generally 3-10m high with many branches and crooked stems (Fig 1). Leaves (5-15cm × 4-6 cm) are opposite, simple, stipules absent, oblong – elliptic, dull grey to yellow green with entire margin, obtuse to bluntly acuminate apex and rounded to subcuneate base with short petiole (Fig 2). Flowers



Figure 1: habit of *P.guajava* L



Figure 2: Dorsal AND Ventral view of the leaves of *P.guajava*-LUCKNOW- 46




Figure 3: T.S through MIDRIB of *P.guajava* L. leaves (LUCKNOW-46)




Figure 4: T.S through MIDRIB of *P.guajava* L. leaves - A portion enlarged- LUCKOW- 46




Figure 5: T.S of LAMINA (LUCKOW- 46)

are white, borne singly or in small clusters, 2-3 cm wide, with 4 or 5 white petals which are quickly shed, and a prominent tuft of perhaps 250 white Stamens . Fruit is small, 3 to 6 cm long, pear-shaped, reddish-yellow when ripe.

#### Microscopy of the leaf:

Transverse section (T.S) of the leaves through the midrib showed the following tissue systems.

Shape: Leaves are dorsiventral with prominent midrib, oblong to elliptic, acuminate. 1.3mm thick, widely concave adaxial side and tangentially oblong abaxial side and semi circular. Lamina is laterally spreading (Fig 3).

Vascular bundle: Broadly bowl shaped vertical- 700µm and 1.6mm horizontal. Xylem is fairly wide thin walled angular in outline occur in compact radial lines. Metaxylem - 30µm wide. Phloem - 50µm wide, encircled around the xylem arc (Fig 4).

Lateral vein: Vascular strands of lateral veins vertically stretched, collateral with sclerenchyma bundle caps.

Mesophyll: Three layers of narrow compact cylindrical palisade cells. Spongy parenchyma is not distinct; two layered vertically elongated compact cells. Tannin content is abundant. Calcium oxalate druses are frequent in the dilated cells.

Ground tissue: Parenchymatous, thin walled compact and tannin content not evident. Secretory cavities occur in the outer zone of adaxial midrib. Calcium oxalate druses are common with dilated ground cells.

Epidermis: 230µm thick, smooth uniformly even. Upper epidermal cells consist of oblong, spindle shaped outer layer and 2 to 3 layers of wider rectangular cells (80- 100µm). Lower epidermis is thin with elliptic narrow cells (15µm thick) (Fig 5).

Powder microscopy: The analysis of the dried powder of the leaf showed paracytic stomata, calcium oxalate druses, circular secretory cavities, xylem vessels, phloem, and fibers and tannin containing cells.

### Physicochemical analysis

Physicochemical parameters were found as follows: total ash 11.05%w/w, acid insoluble ash 1.51, water soluble ash 2.75%w/w, ethanol soluble extractive value 17.98%w/w, water soluble extractive value 20.46%w/w, petroleum ether soluble extractive 2.61%, benzene soluble extractive 4.42%w/w, ethyl acetate soluble extractive 5.61%w/w, chloroform soluble extractive 5.2%w/w, loss on drying 9.7%w/w and foreign organic matter was nil. Leaf constants were as follows vein islet number 3.5, vein termination number 4.3, stomatal number (lower epidermis) 42.4, stomatal number (upper epidermis) 36.2, stomatal index (lower epidermis) 19.3, stomatal index (upper epidermis) 20.1.

### Preliminary phytochemical screening

Preliminary phytochemical screening showed the presence of flavonoids, terpenoids, sterols, tannin, volatile oil, saponins, proteins and amino acids, carbohydrates, reducing sugars, and absence of alkaloids, cyanogenetic glycosides, anthroquinone glycosides, cardiac glycosides, mucilage and fixed oil.

### DISCUSSION

Sensory evaluation plays a key role in determining the suitability or denunciation of a crude drug. Organoleptic testing of a crude drug is mainly for qualitative evaluation based on the observation of morphological and sensory profile. In this report, various morphological, microscopical, physicochemical standards have been developed. Hence we have undertaken this study to serve as a tool for developing standards for identification, quality and purity of

*P.guajava* leaves.

Adulteration and misidentification of crude drugs can cause serious health problems to consumers and legal problems for the pharmaceutical industries. It can be conducted via a variety of techniques, namely macro and microscopic identification and chemical analysis especially description of microscopic botanical aspects to determine definitively the proper species of plant material while it is still in its non extracted form. The observation of cellular level morphology or anatomy is a major aid for the authentication of drugs. These characters are especially important for identification of powdered drugs, because in these cases most of the morphological diagnostic features are lost [28]. Microscopic evaluation is one of the simplest and cheapest methods for the correct identification of the source of the materials [34]. The macroscopic and organoleptic characters of the leaf can serve as diagnostic parameters [35]. Broadly bowl shaped vascular bundle, Xylem is fairly wide thin walled angular in outline occur in compact radial lines. Vascular strands of lateral veins vertically stretched, collateral with sclerenchyma bundle caps. Calcium oxalate druses are common with dilated ground cells were seen.

The ash values are particularly important to find out the presence or absence of foreign inorganic matter such as metallic salts and or silica (earthy matter) [36]. Acid insoluble ash provides information about non-physiological ash produced due do adherence of

inorganic dirt, dust to the crude drug. Increased acid insoluble ash indicates adulteration due do dirt, sand (or) soil. The extractive values are primarily useful for the determination of exhausted or adulterated drug and helpful in the detection of adulteration [37]. Phytochemical evaluation and molecular characterization of plants is an important task in medicinal botany and drug discovery [38]. Preliminary phytochemical screening showed the presence of sterols, flavonoids, terpenoids, saponins, volatile oil, protein and aminoacids, reducing sugars, carbohydrates, and absence of alkaloids, fixed oil, mucilage and glycosides. Dried powder of the leaf showed paracytic stomata, calcium oxalate druses, three layers of wide rectangular cells, circular secretory cavities, broadly bowl shaped vascular bundles, xylem vessels and phloem.

### CONCLUSION

The study of Pharmacognostical features of *Psidium guajava* Linn. (Lucknow-46) had shown the standards which will be useful for the detection of its identity and authenticity. The other study viz. physical evaluation, preliminary phytochemical test add to its quality control and quality assurance for proper identification.

### Conflict of interest statement

We declare that we have no conflict of interest.

### Acknowledgement

The author thanking for all helping hands particularly **Dr.P.Jayaraman**, Director of Plant Anatomy Research Institute,Tambaram, Chennai for microscopical studies.

### REFERENCES

1. Anonymous. *Wealth of India* Raw materials. Vol II. National Institute of Science Communication and Information Resources (NISCAIR), CSIR: New Delhi; 2005,293.
2. Warriar PK, Nambiar VPK, Ramankutty C. *Indian Medicinal Plants – a Compendium of 500 Species*. Vol. IV. Arya Vaidya Sala, Orient Longman Limited: Kottakal; 1995, 371.
3. Nadkarni KM, Nadkarni AK. *Indian Materia Medica*. Vol. I. 3<sup>rd</sup> ed, Popular Prakasham: Mumbai; 1976, 1019.
4. Nundkumar N, Ojewole JA. Studies on the antiparasitodal properties of some South African medicinal plants used as antimalarial remedies in Zulu folk medicine. *Methods Find Exp Clin Pharmacol* 2002; 24(7):397-401.
5. Hernandez, Dolores F. *Plants of the Philippines*. 2<sup>nd</sup> ed. M&L Licudine Enterprises: Phillipines; 1980.
6. Gnan SO, Demello MT. Inhibition of *Staphylococcus aureus* by aqueous *Goiaba* extracts. *J Ethno pharmacol* 1999; 68:103-8.
7. Oliver-Bever, Bep. *Medicinal Plants in tropical West Africa*. Cambridge University Press: Cambridge; 1986.
8. Jaiarj P, Khoohaswan P, Wongkrajang Y, Peungvicha P, Suriyawong P, Saraya M.L.S. Anticough and antimicrobial activities of *Psidium guajava* Linn. leaf extract. *J Ethno pharmacol* 1999; 67(2): 203-12.
9. Dutta BK, Rahman I, Das TK. *In vitro* study on antifungal property of common fruit plants. *Biomedicine* 2000; 20(3): 187-89.
10. Lin J, Puckree T, Mvelase TP. Anti-diarrhoeal evaluation of some medicinal plants used by Zulu traditional healers. *J Ethno pharmacol* 2002; 79(1): 53-6.
11. Ticzon, Romeo. *Ticzon Herbal Medicine Encyclopaedia*. Ticzon Publishing: Phillipines; 1997.
12. Muruganandan S, Srinivasan K, Tandan SK, Jawahar Lal, Chandra S, Raviprakash V. Anti-inflammatory and analgesic activities of some medicinal plants. *Journal of Medicinal and Aromatic Plant Sciences* 2001; 22/23: 56-8.
13. Kavimani S, Karpagam RI, Jaykar B. Anti-inflammatory activity of volatile oil of *Psidium guajava*. *Indian Journal of Pharmaceutical Sciences* 1997; 59(3): 142-44.
14. Conway, Peter. *Tree Medicine – a comprehensive guide to the healing power of over 170 trees*. Judy Piatkus (Publishers) Ltd: 2001.

15. Wyk, Ben-Erik van, Oudtshoorn, Bosch van, Gericke N. Medicinal Plants of South Africa. 1<sup>st</sup> ed. Briza Publications: Pretoria, South Africa; 1997.
16. Jain SK. Dictionary of Indian folk medicine and Ethnobotany. Deep Publications: New Delhi; 1991.
17. Qian H, Nihorimbere V. Antioxidant power of phytochemicals from *Psidium guajava* leaf. J Zheji Univ Sci 2004; 5(6): 676-83.
18. Ojewole JA. Hypoglycemic and hypotensive effects of *Psidium guajava* Linn leaf aqueous extract. Methods Find Exp Clin Pharmacol 2005; 27(10): 689-95.
19. Ojewole JA. Anti inflammatory and analgesic effects of *Psidium guajava* Linn leaf aqueous extract in rats and mice. Methods Find Exp Clin Pharmacol 2006; 28(7): 441-46.
20. Roy CK, Kamath JV, Asad M. Hepatoprotective activity of *Psidium guajava* Linn leaf extract. Indian J Exp Biol 2006; 44(4): 305-11.
21. Hsieh CL, Lin YC, Yen GC, Chen HY. Preventive effects of guava (*Psidium guajava*) leaves and its active compound against  $\alpha$ -dicarbonyl compounds induced blood coagulation. Food chemistry 2007; 103(2): 528-35.
22. Lozoya X, Meckes M, Abouzaid M, Tortoriello J, Nozzolillo, Arnason JT. Quercetin glycosides in *Psidium guajava* L leaves and determination of spasmolytic principle. Arch Med Res 1994; 25(1): 11-15.
23. Begum S, Hassan SI, Siddiqui BS, Shaheen F, Ghayur MN, Gilani AH. Triterpenoids from the leaves of *Psidium guajava*. Phytochemistry 2002; 61(4): 399-403.
24. Kokate CK, Gokhale SB, Purohit AP. Pharmacognosy. 32<sup>nd</sup> ed. Nirali Prakashan: New Delhi; 2005.111-13.
25. Asokan J. Botanical microtechnique principles and practice.1<sup>st</sup> edn. Plant anatomy research centre: Chennai; 2006.
26. Johansen DA. Plant Microtechnique. MC Graw hill: Newyork; 1940, 523.
27. Evan WC. Trease and Evans Pharmacognosy. 15<sup>th</sup> edn. Elsevier: London Saunders; 2002. 544.
28. WHO. Quality control methods for medicinal plant materials. World Health Organisation: Geneva; 1998.
29. Anonymous. The Ayurvedic Pharmacopoeia of India. Part – I, Vol 11, 1<sup>st</sup> ed. Ministry of Health and family welfare, Department of Indian system of medicine and Homeopathy: New Delhi; 1999, 142-145.
30. Anonymous. The Indian Pharmacopoeia. Vol 11. Ministry of Health and family welfare: New Delhi; 1996, A- 53, 54, 89.
31. Wallis TE. Text book of Pharmacognosy. 5<sup>th</sup> ed. CBS Publishers and Distributors: New Delhi; 1997. 111-17, 561-63
32. Mukherjee PK. Quality control of Herbal drugs- An approach to evaluation of botanicals 1<sup>st</sup> edn. Business Horizon: New Delhi; 2002.
33. Kokate CK, Purohit AP, Gokhale SB. Practical Pharmacognosy. 4<sup>th</sup> ed. Vallabh prakashan: New Delhi; 1997. 117-19, 123-25.
34. Patel S, Zaveri M. Pharmacognostic study of the root of *Justicia gendarussa* Burm. J Trad Med 2011; 6(2): 61-72.
35. Singh S, Machawal L, Chauhan MG. Pharmacognostic study of male leaves of *Trichosanthes dioica* Roxb. with special emphasis on microscopic technique. J Pharmacognosy Phytother 2010; 2(5): 71-75.
36. Nayak BS, Patel KN. Pharmacognostic studies of the *Jatropha curcas* leaves. Int J Pharm Tech Res 2010; 2(1): 140-143.
37. Thomas S, Patil DA, Patil AG, Chandra N. Pharmacognostic evaluation and physicochemical analysis of *Averrhoa carambola* L. fruit. J Herb Toxicol 2008; 2(2): 51-54.
38. Yashvanth S, Rani SS, Rao AS, Madhavendra SS. Anatomical exploration of *Leucas aspera* (Willd) links a medicinal herb and its pharmacognostic relevance. J Pharma Res 2011; 4(12):4777-79.