



CONCLUSION

The analysis of X-ray diffraction patterns revealed the formation of single-phase cubic spinel structure of all the samples under investigation. The cation distribution data suggest that Ti ions occupy octahedral B site, Zn ions goes to tetrahedral A site, Mg and Fe ions prefers to occupy both tetrahedral A and octahedral B site.

REFERENCES

1. R. Bueno, M. L. Gregori, M. C. S. Bobrega, J. Mater. Chem. Phys. (2007)
2. N. Reslescu, E. Rezlescu, C. Sava, F. Tudorache, P. Popa, Phys. Stat. Sol. (a) 201 (2004) 17.
3. Mansour Al-Haj, J. Magn. Magn. Mater. 299 (2006) 435
4. M. A. Ahmed, G. Abd-Ellatif, M. Rashad, J. Magn. Magn. Mater. 232 (2001) 194
5. K. Isawa, J. Sugiyama, H. Yyamauchi, Phys. Rev. B 49 (2) (1994) 1987
6. J. B. Goodenough, A. L. Leob, Phys. Rev. 98 (2) (1955) 391.
7. F. Scholl, K. Binder, Z. Phys. B. 39 (1980) 239
8. Y. Purusham, V. D. Reddy, D. R. Sogar, P. Kishan, P. Venugopal Reddy, Phys. Stat. Sol. (a) 140 (1993) k29.
9. J. Smit, H. Wijn, Ferrites, Philips Technical Library, Cleaver-Home Press Ltd. The Netherlands 1959, p. 299.
10. G. Iancu, I. Mirebeau, M. Hennion, G. gaviolle, J. Hubsch Phys. Re. B 54 (1966) 22.
11. Y. Abbas, M. A. Ahed, M. A. Semary, J. Mater. Sci. 18 (1983) 2890.
12. H. H. Joshi, R. G. Kulkarni, J. Mater. Sci. 21 (1986) 2138.